



HAL
open science

Online estimation of Hilbert-Schmidt operators and application to kernel reconstruction of neural fields

Lucas Brivadis, Antoine Chaillet, Jean Auriol

► **To cite this version:**

Lucas Brivadis, Antoine Chaillet, Jean Auriol. Online estimation of Hilbert-Schmidt operators and application to kernel reconstruction of neural fields. 2022. hal-03660185

HAL Id: hal-03660185

<https://hal-centralesupelec.archives-ouvertes.fr/hal-03660185>

Preprint submitted on 5 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Online estimation of Hilbert-Schmidt operators and application to kernel reconstruction of neural fields

Lucas Brivadis¹, Antoine Chaillet¹, and Jean Auriol¹

¹Université Paris-Saclay, CNRS, CentraleSupélec, Laboratoire des Signaux et Systèmes, 91190, Gif-sur-Yvette, France. Emails: `lucas.brivadis@centralesupelec.fr`, `antoine.chaillet@centralesupelec.fr`, `jean.auriol@centralesupelec.fr`

May 5, 2022

Abstract

An adaptive observer is designed for online estimation of Hilbert-Schmidt operators from online measurement of the state for some class of nonlinear infinite-dimensional dynamical systems. Convergence is ensured under detectability and persistency of excitation assumptions. The class of systems considered is motivated by an application to kernel reconstruction of neural fields, commonly used to model spatiotemporal activity of neuronal populations. Numerical simulations confirm the relevance of the approach.

1 Introduction

The problem of online estimation of unknown parameters in dynamical systems from measured state variables is a major issue in many control systems. It can be addressed by means of adaptive observers, that are observers estimating the unmeasured part of the state and the unknown parameters simultaneously. The theory of adaptive observer design, well-known for linear finite-dimensional systems (see, e.g., [19]), is still an active area of research when it comes to nonlinear [3, 4, 18] and/or infinite-dimensional [9, 11, 12] systems.

In this paper, we design an adaptive observer for a class of nonlinear infinite-dimensional systems that allows the reconstruction of unknown linear operators appearing in the dynamics. These operators are estimated in the Hilbert-Schmidt topology. Therefore, not only the state of the system

is infinite-dimensional, but also the “parameters” (now, operators) to be estimated.

The specific class of systems we consider is motivated by an application to kernel reconstruction in neural fields. The offline estimation of these kernels is now a classical issue in inverse problems for neuroscience (see [1, 17] and references therein), that can be addressed for instance using a Tikhonov regularization. We instead rely on adaptive observer strategies to address the online estimation problem. The crucial additional constraint is that the reconstruction can only be based on past values of the measurements and estimates. The recent work [6] considers a similar problem but uses finite-dimensional conductance-based models (which differs from the infinite-dimensional Wilson- Cowan type equation considered here), and estimate finite-dimensional parameters (while we reconstruct linear operators on infinite- dimensional spaces).

Organization of the paper The class of infinite-dimensional nonlinear systems under consideration and a precise formulation of the estimation problem are given in Section 2. In Section 3, we show how this problem may be applied to the issue of online reconstruction of kernels in neural fields dynamics. An adaptive observer is proposed in Section 4, and its convergence is proved under a persistency of excitation condition. Finally, numerical simulations are provided in Section 5.

Notation Given a Hilbert space X , we denote by $\langle \cdot, \cdot \rangle_X$ and $\| \cdot \|_X$ its corresponding scalar product and norm. The identity operator over X is denoted by Id_X . If Y is a Hilbert space, we denote by $\mathcal{L}(X, Y)$ the space of bounded linear operators from X to Y . For all $B \in \mathcal{L}(X, Y)$, we denote by $B^* \in \mathcal{L}(Y, X)$ its adjoint. If $B \in \mathcal{L}(X, X)$, we denote by $\text{Tr}(B)$ the trace of B if it exists. For any open interval $I \subset \mathbb{R}$, $L^p(I, X)$ and $W^{m,p}(I, X)$ stand for the usual Lebesgue and Sobolev spaces, endowed with their canonical norms.

2 Problem statement

2.1 Functional setting

Let $(X, \| \cdot \|_X)$ and $(Y, \| \cdot \|_Y)$ be two separable Hilbert spaces. Consider an infinite-dimensional dynamical system of the following form:

$$\begin{cases} \dot{x} = A_1(x) + \psi(y) + u_1 \\ \dot{y} = A_2(y) + B_1\phi_1(x) + B_2\phi_2(y) + u_2 \end{cases} \quad (1)$$

where (x, y) is the state of the system lying in $X \times Y$, u_1 and u_2 are inputs respectively lying in X and Y , and $A_1 : \mathcal{D}(A_1) \rightarrow X$ and $A_2 : \mathcal{D}(A_2) \rightarrow Y$ are

singled-valued m -dissipative operators (see [16] for a definition), respectively defined on dense subsets $\mathcal{D}(A_1) \subset X$ and $\mathcal{D}(A_2) \subset Y$, such that $A_1(0) = 0$ and $A_2(0) = 0$. The linear operators $B_1 \in \mathcal{L}(X, Y)$ and $B_2 \in \mathcal{L}(Y, Y)$ are bounded, and $\psi : Y \rightarrow X$, $\phi_1 : X \rightarrow X$ and $\phi_2 : Y \rightarrow Y$ are Lipschitz continuous on any bounded set. According to [20, Chapter IV, Proposition 3.1], A_1 and A_2 are generators of nonlinear strongly continuous contraction semigroups over X and Y respectively.

It follows from [20, Chapter IV, Theorems 4.1 and 4.1A] that if u_1 and u_2 are absolutely continuous over \mathbb{R}_+ , then for all $(x_0, y_0) \in \mathcal{D}(A_1) \times \mathcal{D}(A_2)$, there exists $t_{\max} \in (0, +\infty]$ such that (1) admits a unique strong solution $(x, y) : [0, t_{\max}) \rightarrow X \times Y$, i.e., such that $(x(0), y(0)) = (x_0, y_0)$, (x, y) is absolutely continuous, satisfies (1) almost everywhere and lies in $\mathcal{D}(A_1) \times \mathcal{D}(A_2)$. Moreover, if (x, y) is bounded in $X \times Y$ over $[0, t_{\max})$, then $t_{\max} = +\infty$.

2.2 Problem formulation

In this paper, we consider the following online estimation problem.

Problem 2.1 *From the knowledge of A_1 , A_2 , ψ , ϕ_1 , ϕ_2 and the online measurement of u_1 , u_2 and y , estimate online x and the operators B_1 and B_2 .*

In addition to the hypotheses made to ensure the well-posedness of the system, we consider the following two main assumptions.

Assumption 2.2 (Strong dissipativity) *The nonlinear operator A_1 is strongly dissipative, that is, there exists a positive constant α such that for all $(x_1, x_2) \in \mathcal{D}(A_1)^2$,*

$$\langle A_1(x_1) - A_1(x_2), x_1 - x_2 \rangle_X \leq -\alpha \|x_1 - x_2\|_X^2. \quad (2)$$

Since A_1 is supposed to be m -dissipative, we already have that $\langle A_1(x_1) - A_1(x_2), x_1 - x_2 \rangle_X \leq 0$, so that Assumption 2.2 is indeed a stronger dissipativity assumption. Assumption 2.2 implies that every solutions of $\dot{x} = A_1(x)$ are exponentially converging to one another in X at exponential rate α . Since y is supposed to be known online while x is unknown, Assumption 2.2 can be interpreted as a detectability hypothesis: the unknown part of the state has a contracting dynamics. In the estimation strategy, this allows to estimate x online simply by simulating a particular trajectory of the x -subsystem (as all other solutions will eventually converge to it). Note that the unmeasured state x can also be assumed of null dimension, in which case the system (1) reduces to

$$\dot{y} = A_2(y) + B_2\phi_2(y) + u_2.$$

All the results of the paper are still valid in that easier case.

Assumption 2.3 (Hilbert-Schmidt operators) *The linear bounded operators B_1 and B_2 are Hilbert-Schmidt operators, that is, for any Hilbert basis $(e_k^1)_{k \in \mathbb{N}}$ and $(e_k^2)_{k \in \mathbb{N}}$ of X and Y respectively,*

$$\begin{aligned}\|B_1\|_{\mathcal{L}_2(X,Y)}^2 &:= \sum_{k \in \mathbb{N}} \|B_1 e_k^1\|_X^2 = \text{Tr}(B_1^* B_1) < +\infty, \\ \|B_2\|_{\mathcal{L}_2(Y,Y)}^2 &:= \sum_{k \in \mathbb{N}} \|B_2 e_k^2\|_X^2 = \text{Tr}(B_2^* B_2) < +\infty,\end{aligned}$$

where B_i^* denotes the adjoint operator of B_i for $i \in \{1, 2\}$.

We denote by $(\mathcal{L}_2(X, Y), \|\cdot\|_{\mathcal{L}_2(X,Y)})$ and $(\mathcal{L}_2(Y, Y), \|\cdot\|_{\mathcal{L}_2(Y,Y)})$ the Hilbert spaces of Hilbert-Schmidt operators from X to Y and Y to Y , respectively. Then, Problem 2.1 consists in finding $\hat{B}_1(t)$ and $\hat{B}_2(t)$ for all $t \geq 0$ such that $\|\hat{B}_1(t) - B_1\|_{\mathcal{L}_2(X,Y)} \rightarrow 0$ and $\|\hat{B}_2(t) - B_2\|_{\mathcal{L}_2(Y,Y)} \rightarrow 0$ as $t \rightarrow +\infty$. Such estimators must only depend on the knowledge of $A_1, A_2, \psi, \phi_1, \phi_2, u_1(s), u_2(s)$ and $y(s)$ for $s \in [0, t]$. Note that the topology induced by the Hilbert-Schmidt norm is finer than the one induced by the operator norm since $\|\cdot\|_{\mathcal{L}} \leq \|\cdot\|_{\mathcal{L}_2}$.

3 Kernel reconstruction of neural fields

3.1 Neural fields

Problem 2.1 is motivated by an application to kernel reconstruction of neural fields. Neural fields are nonlinear integro-differential equations modeling the spatiotemporal evolution of the activity of neuronal populations. They are based on the seminal works [2, 23] and surveys on their extensive use in mathematical neuroscience can be found in [5, 8]. Given a compact set $\Omega \subset \mathbb{R}^q$ (where, typically, $q \in \{1, 2, 3\}$) representing the physical support of the population, the evolution of neuronal activity $z(t, r) \in \mathbb{R}^n$ at time $t \in \mathbb{R}_+$ and position $r \in \Omega$ is modeled as

$$\begin{aligned}\tau(r) \frac{\partial z}{\partial t}(t, r) &= -z(t, r) + \int_{r' \in \Omega} w(r, r') S(z(t, r')) dr' \\ &\quad + u(t, r),\end{aligned}\quad (3)$$

where $\tau(r)$ is a positive diagonal matrix of size $n \times n$ representing the time decay constant of neuronal activity at position r , $S : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a nonlinear activation function (typically, a sigmoid), $w(r, r') \in \mathbb{R}^{n \times n}$ defines a kernel describing the synaptic strength between location r and r' and $u(t, r) \in \mathbb{R}^n$ is an input. We consider the problem of online reconstruction of the kernel w from the measurement of the neuronal activity z .

3.2 Application

Now we show how (3) fits into (1) and discuss the relevance of Assumptions 2.2 and 2.3 in this context. In order to ensure well-posedness, we make the following usual assumptions on S and w :

- S is bounded, differentiable and has bounded derivative;
- w is square-integrable over Ω^2 .

These assumptions are standard in neural fields analysis. In particular, the boundedness of S reflects the biological limitations of the maximal activity that can be reached by the population.

We assume that the neuronal population can be decomposed into $z(t, r) = (z_1(t, r), z_2(t, r)) \in \mathbb{R}^{n-m} \times \mathbb{R}^m$ where z_1 corresponds to the unmeasured part of the state and z_2 to the measured part. Such a decomposition is natural when the two considered populations are physically separated, as it happens in the brain structures involved in Parkinson's disease [7]. It can also be relevant for imagery techniques that discriminate among neuron types within a given population. Accordingly, we define τ_i , S_i , w_{ij} and u_i of suitable dimensions for each population $i, j \in \{1, 2\}$ so that

$$\begin{aligned} \tau_i(r) \frac{\partial z_i}{\partial t}(t, r) = & -z_i(t, r) \\ & + \sum_{j=1}^2 \int_{r' \in \Omega} w_{ij}(r, r') S_j(z_j(t, r')) dr' + u_i(t, r). \end{aligned} \quad (4)$$

Denote by m the dimension of the measured activity $z_2(r, t)$. In order to fit (4) in the form of (1), set $X = L^2(\Omega; \mathbb{R}^{n-m})$, $Y = L^2(\Omega; \mathbb{R}^m)$, $x = \tau_1 z_1$, $y = \tau_2 z_2$, $W_{ij}(z_j) = \int_{r' \in \Omega} w_{ij}(\cdot, r') z_j(r') dr'$, $A_1 = (-\text{Id}_X + W_{11} S_1)$, $\mathcal{D}(A_1) = X$, $A_2 = -\text{Id}_Y$, $\mathcal{D}(A_2) = Y$, $\psi = W_{12} S_2$, $\phi_i = S_i$ and $B_i = W_{2i}$.

Since w is square-integrable, B_1 and B_2 are Hilbert-Schmidt integral operators with kernels w_{21} and w_{22} , hence Assumption 2.3 is satisfied. In order to satisfy the detectability Assumption 2.2, we need to assume that z_1 has a strongly dissipative dynamics, namely, that A_1 is strongly dissipative. Remark that due to the structure of A_1 , this is the case if

$$\ell_1 \|W_{11}\|_{\mathcal{L}(X, X)} < 1 \quad (5)$$

where ℓ_1 is the Lipschitz constant of S_1 . Indeed,

$$\begin{aligned} & \langle A_1(x_1) - A_1(x_2), x_1 - x_2 \rangle_X \\ & = -\|x_1 - x_2\|_X^2 + \langle W_{11}(S_1(x_1) - S_1(x_2)), x_1 - x_2 \rangle_X \\ & \leq -\alpha \|x_1 - x_2\|_X^2 \end{aligned}$$

for $\alpha = 1 - \ell_1 \|W_{11}\|_{\mathcal{L}(X,X)}$. We stress that condition (5) is commonly used in the stability analysis of neural fields [15] and ensures dissipativity even in the presence of axonal propagation delays [13].

We thus assume that each population is either measured online (taken into account in z_2) or unmeasured but stable and with known kernels (taken into account in z_1). Problem 2.1 is now equivalent to online reconstruction of w_{21} and w_{22} (in $L^2(\Omega^2)$) from the online measurement of z_2 and u_i and the knowledge of τ_i, w_{1i}, S_i for all $i \in \{1, 2\}$. Note that if the full state z is measured (i.e. $m = n$), then no dissipative part z_1 of the system is required, hence the full kernel w is to be estimated.

4 Online estimation of hilbert-schmidt operators

4.1 Adaptive observer design

In order to solve Problem 2.1, we propose to consider B_1 and B_2 as additional constant variables to system (1), so that the resulting state space is the Hilbert space $H := X \times Y \times \mathcal{L}_2(X, Y) \times \mathcal{L}_2(Y, Y)$. Set also $\mathcal{D} := \mathcal{D}(A_1) \times \mathcal{D}(A_2) \times \mathcal{L}_2(X, Y) \times \mathcal{L}_2(Y, Y) \subset H$. Inspired by the estimator proposed in [3] for finite-dimensional nonlinear systems, we consider the following observer over H :

$$\begin{cases} \dot{\hat{x}} = A_1(\hat{x}) + \psi(y) + u_1 \\ \dot{\hat{y}} = A_2(y) + \hat{B}_1\phi_1(\hat{x}) + \hat{B}_2\phi_2(y) + u_2 - \beta(\hat{y} - y) \\ \dot{\hat{B}}_1 = -\gamma_1(\hat{y} - y)\phi_1(\hat{x})^* \\ \dot{\hat{B}}_2 = -\gamma_2(\hat{y} - y)\phi_2(y)^* \end{cases} \quad (6)$$

where β, γ_1 , and γ_2 are positive constants, called observer gains, that need to be appropriately tuned to guarantee the convergence of the observer state to the real state. Note that for any v in Y and any w in X (resp. in Y), vw^* lies in $\mathcal{L}_2(X, Y)$ (resp. in $\mathcal{L}_2(Y, Y)$) and $\|vw^*\|_{\mathcal{L}_2(X, Y)} = \|v\|_Y \|w\|_X$ (resp. $\|vw^*\|_{\mathcal{L}_2(Y, Y)} = \|v\|_Y \|w\|_Y$). Reasoning as in Section 2.1, one can show that the cascade system (1)-(6) is well-posed.

4.2 Main result

Our main result, proved in Section 4.3, relies on the notion of persistence of excitation.

Definition 4.1 (Persistence of excitation) *A signal $g : \mathbb{R}_+ \rightarrow V$ is persistently exciting over the Hilbert space V if there exists positive constants T and κ such that*

$$\int_t^{t+T} g(\tau)g^*(\tau)d\tau \geq \kappa \text{Id}_V, \quad \forall t \geq 0. \quad (7)$$

We now provide sufficient conditions for the convergence of the observer (6) to the state of system (1), thus solving Problem 2.1.

Theorem 4.2 (Observer convergence) *Suppose that Assumptions 2.2 and 2.3 are satisfied. Assume moreover that the functions ϕ_1 and ϕ_2 are bounded and that ϕ_1 is globally Lipschitz continuous with constant ℓ_1 . Pick the observer gains $\beta, \gamma_1, \gamma_2$ such that $\gamma_1, \gamma_2 > 0$ and*

$$4\alpha\beta > \ell_1^2 \|B_1\|_{\mathcal{L}(X,Y)}^2. \quad (8)$$

Then, for any absolutely continuous u_1 and u_2 and any solution of (1) defined over \mathbb{R}_+ , any solution of (6) satisfies

$$\lim_{t \rightarrow +\infty} \|\hat{x}(t) - x(t)\|_X = 0, \quad \lim_{t \rightarrow +\infty} \|\hat{y}(t) - y(t)\|_Y = 0,$$

and $\|\hat{B}_1 - B_1\|_{\mathcal{L}_2(X,Y)}$ and $\|\hat{B}_2 - B_2\|_{\mathcal{L}_2(X,Y)}$ remain bounded.

Moreover, if $t \mapsto (\phi_1(x(t)), \phi_2(y(t)))$ is persistently exciting over $X \times Y$ and if ϕ_1 and ϕ_2 are differentiable with bounded derivatives, then

$$\lim_{t \rightarrow +\infty} \|\hat{B}_i(t) - B_i\|_{\mathcal{L}_2(X,Y)} = 0, \quad \forall i \in \{1, 2\}.$$

It is worth noting that the observer gains γ_1 and γ_2 play no qualitative role in the observer convergence. Also, β can always be picked sufficiently large to fulfill (8). The main requirement therefore lies in the persistence of excitation requirement, which is a common hypothesis to ensure convergence of adaptive observers (see for instance [3, 14, 19] in the finite-dimensional context and [10, 12] in the infinite-dimensional case). Roughly speaking, it states that parameters to be estimated are sufficiently “excited” by the system dynamics. However, this assumption is difficult to check in practice since it depends on the trajectories of the system itself. In Section 5, we choose in numerical simulations a persistently exciting input (u_1, u_2) in order to generate persistence of excitation in the signal $(\phi_1(x), \phi_2(y))$. This strategy seems to be numerically efficient, but the theoretical analysis of the link between the persistence of excitation of (u_1, u_2) and $(\phi_1(x), \phi_2(y))$ remains an open question, not only in the present work but also for general classes of adaptive observers.

As developed in Section 3.2, Theorem 4.2 directly applies to the neural fields context. With the notations of Section 3.2, the adaptive observer takes the form

$$\begin{cases} \tau_1 \dot{\hat{z}}_1 = -\hat{z}_1 + W_{11}S_1(\hat{z}_1) + W_{12}S_2(z_2) + u_1 \\ \tau_2 \dot{\hat{z}}_2 = -z_2 + \hat{W}_{21}S_1(\hat{z}_1) + \hat{W}_{22}S_2(z_2) + u_2 \\ \quad \quad \quad \quad \quad \quad \quad \quad \quad - \beta(\hat{z}_2 - z_2) \\ \dot{\hat{W}}_{21} = -\gamma_1(\hat{z}_2 - z_2)S_1(\hat{z}_1)^* \\ \dot{\hat{W}}_{22} = -\gamma_2(\hat{z}_2 - z_2)S_2(z_2)^*. \end{cases} \quad (9)$$

Then, we have the following, which immediately follows from Theorem 4.2 for this particular system.

Corollary 4.3 (Neural fields estimation) *Suppose that S_1 (resp. S_2) is bounded, differentiable, and that its derivative is bounded by some $\ell_1 > 0$ (resp. $\ell_2 > 0$), and that w is square-integrable over Ω^2 . Assuming that (5) is satisfied, pick the observer gains in such a way that $\gamma_1, \gamma_2 > 0$ and*

$$4(1 - \ell_1 \|W_{11}\|_{\mathcal{L}(X,X)})\beta > \ell_1^2 \|B_1\|_{\mathcal{L}(X,Y)}^2. \quad (10)$$

Consider any solution of (4) defined over \mathbb{R}_+ for some absolutely continuous inputs u_1 and u_2 . Then any solution of (9) satisfies

$$\lim_{t \rightarrow +\infty} \|\hat{z}_1(t) - z_1(t)\|_X = 0, \quad \lim_{t \rightarrow +\infty} \|\hat{z}_2(t) - z_2(t)\|_Y = 0$$

and $\|\hat{W}_{21} - W_{21}\|_{\mathcal{L}_2(X,Y)}$ and $\|\hat{W}_{22} - W_{22}\|_{\mathcal{L}_2(X,Y)}$ remain bounded.

Moreover, if $t \mapsto (S_1(z_1(t)), S_2(z_2(t)))$ is persistently exciting over $X \times Y$, then

$$\lim_{t \rightarrow +\infty} \|\hat{W}_{2i}(t) - W_{2i}\|_{\mathcal{L}_2(X,Y)} = 0, \quad \forall i \in \{1, 2\}.$$

Here again, provided that condition (5) holds, β can always be picked large enough to fulfill (10).

4.3 Proof of Theorem 4.2

Consider a solution (x, y) of (1) defined on \mathbb{R}_+ and the corresponding solution $(\hat{x}, \hat{y}, \hat{B}_1, \hat{B}_2)$ of (6). The estimation error $(\tilde{x}, \tilde{y}, \tilde{B}_1, \tilde{B}_2) := (\hat{x}, \hat{y}, \hat{B}_1, \hat{B}_2) - (x, y, B_1, B_2)$ is ruled by:

$$\begin{cases} \dot{\tilde{x}} = A_1(\hat{x}) - A_1(x) \\ \dot{\tilde{y}} = \hat{B}_1 \phi_1(\hat{x}) - B_1 \phi_1(x) + \tilde{B}_2 \phi_2(y) - \beta \tilde{y} \\ \dot{\tilde{B}}_1 = -\gamma_1 \tilde{y} \phi_1(\hat{x})^* \\ \dot{\tilde{B}}_2 = -\gamma_2 \tilde{y} \phi_2(y)^*. \end{cases} \quad (11)$$

4.3.1 Proof that $(\tilde{x}, \tilde{y}) \rightarrow 0$

We endow H with the squared norm $\|\cdot\|_H^2 = \|\cdot\|_X^2 + \|\cdot\|_Y^2 + \frac{1}{\gamma_1} \|\cdot\|_{\mathcal{L}_2(X,Y)}^2 + \frac{1}{\gamma_2} \|\cdot\|_{\mathcal{L}_2(Y,Y)}^2$, which is equivalent to the squared norm induced by the Cartesian product $H = X \times Y \times \mathcal{L}_2(X,Y) \times \mathcal{L}_2(Y,Y)$. Given any initial state, denote by $t_{\max} \in (0, +\infty]$ the maximal time of existence of $(\tilde{x}, \tilde{y}, \tilde{B}_1, \tilde{B}_2)$.

Using $\hat{B}_1\phi_1(\hat{x}) - B_1\phi_1(x) = \tilde{B}_1\phi_1(\hat{x}) + B_1(\phi_1(\hat{x}) - \phi_1(x))$, we have almost everywhere on $[0, t_{\max})$

$$\begin{aligned} \frac{d}{dt} \|(\tilde{x}, \tilde{y}, \tilde{B}_1, \tilde{B}_2)\|_H^2 &= \langle A_1(\hat{x}) - A_1(x), \tilde{x} \rangle_X - \beta \|\tilde{y}\|_Y^2 + \langle \tilde{B}_2\phi_2(y), \tilde{y} \rangle_Y \\ &\quad + \langle \tilde{B}_1\phi_1(\hat{x}), \tilde{y} \rangle_Y + \langle B_1(\phi_1(\hat{x}) - \phi_1(x)), \tilde{y} \rangle_Y \\ &\quad - \langle \tilde{y}\phi_1(\hat{x})^*, \tilde{B}_1 \rangle_{\mathcal{L}_2(X,Y)} - \langle \tilde{y}\phi_1(y)^*, \tilde{B}_2 \rangle_{\mathcal{L}_2(X,Y)}. \end{aligned}$$

By Assumption 2.2, $\langle A_1(\hat{x}) - A_1(x), \tilde{x} \rangle_X \leq -\alpha \|\tilde{x}\|_X^2$. By definition of the Hilbert-Schmidt scalar product,

$$\begin{aligned} \langle \tilde{y}\phi_1(\hat{x})^*, \tilde{B}_1 \rangle_{\mathcal{L}_2(X,Y)} &= \text{Tr}(\phi_1(\hat{x})\tilde{y}^*\tilde{B}_1) = \text{Tr}(\tilde{y}^*\tilde{B}_1\phi_1(\hat{x})) \\ &= \langle \tilde{B}_1\phi_1(\hat{x}), \tilde{y} \rangle_Y \end{aligned}$$

and, similarly,

$$\langle \tilde{y}\phi_2(y)^*, \tilde{B}_2 \rangle_{\mathcal{L}_2(X,Y)} = \langle \tilde{B}_2\phi_2(y), \tilde{y} \rangle_Y.$$

Hence

$$\begin{aligned} \frac{d}{dt} \|(\tilde{x}, \tilde{y}, \tilde{B}_1, \tilde{B}_2)\|_H^2 &\leq -\alpha \|\tilde{x}\|_X^2 - \beta \|\tilde{y}\|_Y^2 + \langle B_1(\phi_1(\hat{x}) - \phi_1(x)), \tilde{y} \rangle_X. \end{aligned}$$

By Cauchy-Schwartz inequality, for all $\varepsilon > 0$,

$$\begin{aligned} \langle B_1(\phi_1(\hat{x}) - \phi_1(x)), \tilde{y} \rangle_X &\leq \ell_1 \|B_1\|_{\mathcal{L}(X,Y)} \left(\frac{\varepsilon}{2} \|\tilde{x}\|_X^2 + \frac{1}{2\varepsilon} \|\tilde{y}\|_X^2 \right) \end{aligned}$$

where ℓ_1 is the Lipschitz constant of ϕ_1 . Pick $\varepsilon = \frac{\alpha}{\ell_1 \|B_1\|_{\mathcal{L}(X,Y)}} + \frac{\ell_1 \|B_1\|_{\mathcal{L}(X,Y)}}{4\beta} > 0$. Using condition (8), we get that $\mu_1 := \alpha - \ell_1 \|B_1\|_{\mathcal{L}(X,Y)}\varepsilon/2 > 0$ and $\mu_2 := \beta - \ell_1 \|B_1\|_{\mathcal{L}(X,Y)}/2\varepsilon > 0$. Then

$$\frac{d}{dt} \|(\tilde{x}, \tilde{y}, \tilde{B}_1, \tilde{B}_2)\|_H^2 \leq -\mu_1 \|\tilde{x}\|_X^2 - \mu_2 \|\tilde{y}\|_Y^2.$$

Thus $(\tilde{x}, \tilde{y}, \tilde{B}_1, \tilde{B}_2)$ remains bounded. Hence according to [20, Chapter IV, Theorems 4.1], we obtain $t_{\max} = +\infty$ i.e. the state $(\hat{x}, \hat{y}, \hat{B}_1, \hat{B}_2)$ is defined over \mathbb{R}_+ . Moreover, we have $\frac{d}{dt} \|\tilde{x}\|_X^2 \leq 0$ and

$$\begin{aligned} \frac{d}{dt} \|\tilde{y}\|_Y^2 &\leq -\beta \|\tilde{y}\|_Y^2 + \langle \tilde{B}_2\phi_2(y), \tilde{y} \rangle_Y \\ &\quad + \langle \hat{B}_1\phi_1(\hat{x}), \tilde{y} \rangle_Y - \langle B_1\phi_1(x), \tilde{y} \rangle_Y \end{aligned}$$

which is bounded since $(\tilde{x}, \tilde{y}, \tilde{B}_1, \tilde{B}_2)$ is bounded, B_1 is constant, and ϕ_1 and ϕ_2 are bounded. Hence, according to Barbalat's lemma, $(\tilde{x}(t), \tilde{y}(t)) \rightarrow 0$ as $t \rightarrow +\infty$.

4.3.2 Proof that $(\tilde{B}_1, \tilde{B}_2) \rightarrow 0$

Now assume that $t \mapsto (\phi_1(x(t)), \phi_2(y(t)))$ is persistently exciting over $X \times Y$ and that ϕ_1 and ϕ_2 are differentiable with bounded derivatives. Note that $\hat{B}_1\phi_1(\hat{x}) - B_1\phi_1(x) = \hat{B}_1(\phi_1(\hat{x}) - \phi_1(x)) + \tilde{B}_1\phi_1(x)$. Hence the error dynamics (11) can be written as

$$\begin{cases} \dot{\tilde{y}}(t) = \tilde{B}_1(t)g_1(t) + \tilde{B}_2(t)g_2(t) + f_0(t) \\ \dot{\tilde{B}}_1(t) = f_1(t) \\ \dot{\tilde{B}}_2(t) = f_2(t), \end{cases} \quad (12)$$

where $f_0(t) := \hat{B}_1(\phi_1(\hat{x}(t)) - \phi_1(x(t))) - \beta\tilde{y}$, $f_1(t) := -\gamma_1\tilde{y}(t)\phi_1(\hat{x}(t))^*$, $f_2(t) := -\gamma_2\tilde{y}(t)\phi_2(y(t))^*$, $g_1(t) := \phi_1(x(t))$ and $g_2(t) := \phi_2(y(t))$ for all $t \geq 0$. Since $(\tilde{x}(t), \tilde{y}(t)) \rightarrow 0$ as $t \rightarrow +\infty$, \hat{B}_1 is bounded, ϕ_1 is globally Lipschitz and ϕ_1 and ϕ_2 are bounded, we get that $f_i(t)$ tends toward 0 as t goes to $+\infty$ for all $i \in \{0, 1, 2\}$. Set $g := (g_1, g_2) : \mathbb{R}_+ \rightarrow X \times Y$ and $f_{1,2}, \tilde{B} : \mathbb{R}_+ \rightarrow \mathcal{L}_2(X \times Y, Y)$, defined by $f_{1,2}(t)(\zeta, \xi) = f_1(t)\zeta + f_2(t)\xi$, $\tilde{B}(t)(\zeta, \xi) = \tilde{B}_1(t)\zeta + \tilde{B}_2(t)\xi$ for all $(\zeta, \xi) \in X \times Y$ and all $t \geq 0$, so that $\dot{\tilde{y}}(t) = \tilde{B}(t)g(t) + f_0(t)$. Remark that $\|\tilde{B}(t)\|_{\mathcal{L}_2(X \times Y, Y)}^2 = \|\tilde{B}_1(t)\|_{\mathcal{L}_2(X, Y)}^2 + \|\tilde{B}_2(t)\|_{\mathcal{L}_2(Y, Y)}^2$, so that it remains to show that $\|\tilde{B}(t)\|_{\mathcal{L}_2(X \times Y, Y)} \rightarrow 0$ as $t \rightarrow +\infty$ to conclude. Applying twice Duhamel's formula, we have for all $t, \tau \geq 0$:

$$\begin{aligned} \tilde{y}(t + \tau) &= \tilde{y}(t) + \tilde{B}(t) \int_0^\tau g(t + s) ds \\ &\quad + \int_0^\tau \int_0^s f_{1,2}(t + \sigma) g(t + s) d\sigma ds + \int_0^\tau f_0(t + s) ds. \end{aligned}$$

Define $\mathcal{O}(t, T) := \int_0^T \|\tilde{y}(t + \tau)\|_Y^2 d\tau$ for any $T > 0$ and $t \geq 0$. Since $\tilde{y}(t) \rightarrow 0$, $\mathcal{O}(t, T) \rightarrow 0$ as $t \rightarrow +\infty$. Moreover,

$$\begin{aligned} \mathcal{O}(t, T) &= \int_0^T \left\| \tilde{y}(t) + \int_0^\tau \int_0^s f_{1,2}(t + \sigma) g(t + s) d\sigma ds \right. \\ &\quad \left. + \int_0^\tau f_0(t + s) ds \right\|_Y^2 d\tau + \int_0^T \left\| \tilde{B}(t) \int_0^\tau g(t + s) ds \right\|_Y^2 d\tau \\ &\quad + \int_0^T \left\langle \tilde{y}(t) + \int_0^\tau \int_0^s f_{1,2}(t + \sigma) g(t + s) d\sigma ds \right. \\ &\quad \left. + \int_0^\tau f_0(t + s) ds, \tilde{B}(t) \int_0^\tau g(t + s) ds \right\rangle_Y d\tau. \end{aligned}$$

Since $\tilde{y}(t)$ and $f_i(t)$ tends toward 0 as t goes to $+\infty$ for all $i \in \{0, 1, 2\}$ and g and \tilde{B} are bounded, we get that

$$\lim_{t \rightarrow +\infty} \int_0^T \left\| \tilde{B}(t) \int_0^\tau g(t + s) ds \right\|_Y^2 d\tau = 0. \quad (13)$$

For all $t \geq 0$, define $h(t, \tau) = \tilde{B}(t) \int_0^\tau g(t+s) ds$. By (13), $\|h(t, \cdot)\|_{L^2((0,T);Y)} \rightarrow 0$ as $t \rightarrow +\infty$. Note that $\frac{\partial h}{\partial \tau}(t, \tau) = \tilde{B}(t)g(t+\tau)$ hence $h(t, \cdot) \in W^{1,2}((0,T);Y)$ since g is bounded. Moreover, \dot{y} is bounded since \tilde{B}_i and g_i are bounded for $i \in \{1,2\}$ and $\dot{x} = A_1(\hat{x}) - A_1(x)$ is bounded since A_1 is m -dissipative (see [16, Corollary 3.7 and Theorem 4.20]). Hence, if ϕ_1 and ϕ_2 are differentiable with bounded derivatives, then so is g . Therefore, for all $t \geq 0$, $h(t, \cdot) \in W^{2,2}((0,T);Y)$ and $\|h(t, \cdot)\|_{W^{2,2}((0,T);Y)} \leq C_1$ for some positive constant C_1 independent of t . According to the interpolation inequality (see, e.g., [21, Section II.2.1]),

$$\|h(t, \cdot)\|_{W^{1,2}((0,T);Y)}^2 \leq C_2 \|h(t, \cdot)\|_{L^2((0,T);Y)}$$

for some positive constant C_2 independent of t . Thus $\|\frac{\partial h}{\partial \tau}(t, \tau)\|_{L^2((0,T);Y)} \rightarrow 0$, meaning that

$$\lim_{t \rightarrow +\infty} \int_0^T \left\| \tilde{B}(t)g(t+\tau) \right\|_Y^2 d\tau = 0. \quad (14)$$

Now, let $(e_k)_{k \in \mathbb{N}}$ be a Hilbert basis of Y . Then

$$\begin{aligned} \|\tilde{B}(t)\|_{\mathcal{L}_2(X \times Y, Y)}^2 &= \|\tilde{B}^*(t)\|_{\mathcal{L}_2(Y, X \times Y)}^2 \\ &= \sum_{k \in \mathbb{N}} \|\tilde{B}(t)^* e_k\|_Y^2. \end{aligned}$$

Since g is persistently exciting, we have, for some $T, \kappa > 0$,

$$\int_0^T |\langle g(t+\tau), v \rangle_{X \times Y}|^2 d\tau \geq \kappa \|v\|_{X \times Y}^2, \quad \forall t \geq 0,$$

for all $v \in X \times Y$. Then,

$$\begin{aligned} \kappa \|\tilde{B}(t)\|_{\mathcal{L}_2(X \times Y, Y)}^2 &\leq \sum_{k \in \mathbb{N}} \int_0^T |\langle g(t+\tau), \tilde{B}(t)^* e_k \rangle_{X \times Y}|^2 d\tau \\ &= \sum_{k \in \mathbb{N}} \int_0^T |\langle \tilde{B}(t)g(t+\tau), e_k \rangle_Y|^2 d\tau \\ &= \int_0^T \sum_{k \in \mathbb{N}} |\langle \tilde{B}(t)g(t+\tau), e_k \rangle_Y|^2 d\tau \\ &= \int_0^T \left\| \tilde{B}(t)g(t+\tau) \right\|_Y^2 d\tau. \end{aligned}$$

Thus, by (14), $\|\tilde{B}(t)\|_{\mathcal{L}_2(X \times Y, Y)} \rightarrow 0$ as $t \rightarrow +\infty$, which concludes the proof. \blacksquare

5 Numerical simulation

We provide a numerical simulation of the adaptive observer (6) in the case of a two-dimensional neural field (namely, $n = 2$ and $m = 1$ in Section 3.2) over the unit circle $\Omega = \mathbb{S}^1$. We set parameters of the system and observer as in Table 1, so that all assumptions of Corollary 4.3 are satisfied. Initial conditions are given by $z_1(0) = z_2(0) = 1$, $\hat{z}_1(0) = \hat{z}_2(0) = 0$ and $\hat{B}_1(0) = \hat{B}_2(0) = 0$. Kernels are given by Gaussian functions depending on the distance between r and r' , as it is frequently assumed in practice (see [7]): $w_{ij}(r, r') = \omega_{ij} \mathbf{g}(r, r') / \|\mathbf{g}\|_{L^2(\Omega^2; \mathbb{R})}$, $\mathbf{g}(r, r') = \exp(-\sigma|r - r'|^2)$ for constant parameters σ and ω_{ij} given in Table 1. The inputs u_i are chosen as spatiotemporal periodic signals with irrational frequency ratio, i.e., $u_i(t, r) = 10^3 \sin(\lambda_i t r)$ with λ_1/λ_2 irrational. This choice is made to ensure persistency of excitation of the input (u_1, u_2) , which in practice seems to be sufficient to ensure persistency of excitation of $(S_1(z_1), S_2(z_2))$. Note that for $u_1 = u_2 = 0$, the persistency of excitation assumption seems to be not guaranteed, hence the observer does not converge. However, in practice, such a persistent input is likely to occur due to exogenous signals coming from other unmodeled neuronal populations.

In numerical simulations, the system is spatially discretized over Ω with a constant space step $\Delta r = 1/20$, and the resulting ordinary differential equation is solved with an explicit Runge-Kutta (4, 5) method.

$S_i(z) = \tanh(z)$	$\tau_i = 1$	$\lambda_1 = 1$	$\lambda_2 = \sqrt{2}$
$\omega_{11} = 0.1$	$\omega_{12} = 2$	$\omega_{21} = -2$	$\omega_{22} = 2$
$\beta = 1$	$\gamma_1 = 100$	$\gamma_2 = 100$	$\sigma = 60$

Table 1: System and observer parameters for the numerical simulation of Figures 1 and 2

In Figure 1, the convergence of the observer, that is proved in Corollary 4.3, is numerically verified. In Figure 2, we illustrate some iterations of the reconstructed kernel $\hat{w}_{22}(t)$ of $\hat{B}_2(t)$, which converges to the kernel w_{22} of B_2 .

6 Conclusion

In this paper, we have shown that an observer can be designed to estimate online linear operators arising in some nonlinear infinite-dimensional dynamical systems from the measurement of the state. This estimation problem is motivated by an application to kernel reconstruction for neural fields equations. The main assumption is the persistence of excitation of the system along its trajectories. Our simulations suggest that this requirement can be ensured using appropriate exogenous inputs. In future works, we wish to




Figure 1: Evolution of the estimation errors $\|\hat{B}_1(t) - B_1\|_{\mathcal{L}_2(X,Y)}$, $\|\hat{B}_2(t) - B_2\|_{\mathcal{L}_2(X,Y)}$, $\|\hat{x}(t) - x(t)\|_X$ and $\|\hat{y}(t) - y(t)\|_Y$, for system and observer parameters given in Table 1.

investigate this hypothesis by either designing inputs ensuring persistence, or designing observers that do not rely on this assumption (see, e.g., [22]). Moreover, the use of this estimator in closed-loop to stabilize the systems by means of dynamic output feedback will be investigated. Finally, delayed neural fields in the form of [7] could be considered, as they do not fit into the functional setting of the present paper although capturing meaningful biological processes such as non-instantaneous axonal propagation.

References

- [1] J. Alswaihi, R. Potthast, I. Bojak, D. Saddy, and A. Hutt. Kernel reconstruction for delayed neural field equations. *The Journal of Mathematical Neuroscience*, 8(1):1–24, 2018.
- [2] S. Amari. Dynamics of pattern formation in lateral-inhibition type neural fields. *Biological cybernetics*, 27(2):77–87, 1977.
- [3] G. Besançon. Remarks on nonlinear adaptive observer design. *Systems & Control Letters*, 41(4):271–280, 2000.
- [4] G. Besançon and A. Țiclea. On adaptive observers for systems with state and parameter nonlinearities. *IFAC-PapersOnLine*, 50(1):15416–15421, 2017. 20th IFAC World Congress.




Figure 2: Evolution of the kernel $\hat{w}_{22}(t, r, r')$ of $\hat{B}_2(t)$ at times $t = 0$, $t = 250$, $t = 500$ and $t = 1000$.

- [5] P. Bressloff. Spatiotemporal dynamics of continuum neural fields. *Journal of Physics A: Mathematical and Theoretical*, 45(3), 2012.
- [6] T. B. Burghi and R. Sepulchre. Online estimation of biophysical neural networks, 2021.
- [7] A. Chaillet, G. Detorakis, S. Palfi, and S. Senova. Robust stabilization of delayed neural fields with partial measurement and actuation. *Automatica*, 83:262–274, Sep. 2017.
- [8] S. Coombes, P. beim Graben, R. Potthast, and J. Wright. *Neural Fields: Theory and Applications*. Springer, 2014.
- [9] R. Curtain, M. Demetriou, and K. Ito. Adaptive observers for structurally perturbed infinite dimensional systems. In *Proceedings of the 36th IEEE Conference on Decision and Control*, volume 1, pages 509–514 vol.1, 1997.

- [10] R. Curtain, M. Demetriou, and K. Ito. Adaptive observers for slowly time varying infinite dimensional systems. In *Proceedings of the 37th IEEE Conference on Decision and Control (Cat. No.98CH36171)*, volume 4, pages 4022–4027 vol.4, 1998.
- [11] M. A. Demetriou. Design of adaptive output feedback synchronizing controllers for networked pdes with boundary and in-domain structured perturbations and disturbances. *Automatica*, 90:220–229, 2018.
- [12] M. A. Demetriou and K. Ito. Adaptive observers for a class of infinite dimensional systems. *IFAC Proceedings Volumes*, 29(1):5346–5350, 1996. 13th World Congress of IFAC, 1996, San Francisco USA, 30 June - 5 July.
- [13] G. I. Detorakis and A. Chaillet. Incremental stability of spatiotemporal delayed dynamics and application to neural fields. In *56th IEEE Conference on Decision and Control*, pages 5937–5942, 2017.
- [14] M. Farza, M. M’Saad, T. Maatoug, and M. Kamoun. Adaptive observers for nonlinearly parameterized class of nonlinear systems. *Automatica*, 45(10):2292–2299, 2009.
- [15] O. Faugeras, F. Grimbert, and J.-J. Slotine. Absolute stability and complete synchronization in a class of neural fields models. *SIAM Journal of Applied Mathematics*, 61(1):205–250, 2008.
- [16] I. Miyadera. *Nonlinear semigroups*, volume 109. American Mathematical Soc., 1992.
- [17] R. Potthast and P. B. Graben. Inverse problems in neural field theory. *SIAM Journal on Applied Dynamical Systems*, 8(4):1405–1433, 2009.
- [18] A. Pyrkin, A. Bobtsov, R. Ortega, and A. Isidori. An adaptive observer for uncertain linear time-varying systems with unknown additive perturbations, 2021.
- [19] S. Sastry, M. Bodson, and J. F. Bartram. Adaptive control: stability, convergence, and robustness, 1990.
- [20] R. E. Showalter. *Monotone operators in Banach space and nonlinear partial differential equations*, volume 49. American Mathematical Society, 2013.
- [21] R. Temam. Infinite-dimensional dynamical systems. *Nonlinear functional analysis and its applications, Part 2*, 45(Part 2):431, 1986.
- [22] J. Wang, D. Efimov, and A. A. Bobtsov. On robust parameter estimation in finite-time without persistence of excitation. *IEEE Transactions on Automatic Control*, 65(4):1731–1738, 2020.

- [23] H. R. Wilson and J. D. Cowan. A mathematical theory of the functional dynamics of cortical and thalamic nervous tissue. *Kybernetik*, 13(2):55–80, 1973.