



**HAL**  
open science

# Maximum likelihood estimation and prediction error for a Matérn model on the circle

Sébastien Petit

► **To cite this version:**

Sébastien Petit. Maximum likelihood estimation and prediction error for a Matérn model on the circle. 2022. hal-03778527

**HAL Id: hal-03778527**

**<https://hal-centralesupelec.archives-ouvertes.fr/hal-03778527>**

Preprint submitted on 15 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Maximum likelihood estimation and prediction error for a Matérn model on the circle

Sébastien J. Petit<sup>\*1, 2</sup>

<sup>1</sup> Université Paris-Saclay, CNRS, CentraleSupélec,  
Laboratoire des signaux et systèmes, 91190, Gif-sur-Yvette, France

<sup>2</sup>Laboratoire National de Métrologie et d'Essais, 78197 Trappes Cedex, France

September 15, 2022

## Abstract

This work considers Gaussian process interpolation with a periodized version of the Matérn covariance function (Stein, 1999, Section 6.7) with Fourier coefficients  $\phi(\alpha^2 + j^2)^{-\nu-1/2}$ . Convergence rates are studied for the joint maximum likelihood estimation of  $\nu$  and  $\phi$  when the data is sampled according to the model. The mean integrated squared error is also analyzed with fixed and estimated parameters, showing that maximum likelihood estimation yields asymptotically the same error as if the ground truth was known. Finally, the case where the observed function is a “deterministic” element of a continuous Sobolev space is also considered, suggesting that bounding assumptions on some parameters can lead to different estimates.

## 1 Introduction

Gaussian process regression or kriging is a common technique for inferring an unknown function from data, which has applications in geostatistics (Stein, 1999), computer experiments (Santner et al., 2003), and machine learning (Rasmussen and Williams, 2006). Stein (1999) stresses the importance of choosing a covariance function that fits the problem, promotes the use of the Matérn (1986) family of covariance functions, and advocates using maximum likelihood to estimate its parameters.

A distinction is generally made between increasing and fixed-domain asymptotic frameworks for parameter estimation of Gaussian processes (see, e.g., Bachoc, 2021, for a review). While several increasing-domain asymptotic frameworks have been exhaustively studied (see, e.g., Mardia and Marshall, 1984; Bachoc, 2014), fixed-domain frameworks are studied only with simplifications and for a restrained number of parameters to our knowledge (Ying, 1991, 1993; van der Vaart, 1996; Zhang, 2004; Loh, 2006; Anderes, 2010).

---

<sup>\*</sup>sebastien.petit@lne.fr

Considering fixed-domain asymptotics, the regularity parameter of the Matérn covariance function seems to have been little studied, although [Stein \(1999\)](#) presents numerous results suggesting it as the most impactful from the point of view of prediction error. To study parameter estimation, [Stein \(1999, Section 6.7\)](#) proposes an asymptotic framework<sup>1</sup> with uniformly distributed observations on the torus and makes a conjecture about the asymptotic behavior of maximum likelihood based on the Fisher information matrix. This topic has only recently regained popularity. Indeed, [Chen et al. \(2021\)](#) used the previous framework to show that the estimation of the regularity parameter is consistent if the other parameters remain fixed. Moreover, [Karvonen \(2022\)](#) has recently shown an asymptotic lower bound in the general case of a “nice” bounded domain of  $\mathbb{R}^d$ , also covering the case of a deterministic function from a continuous Sobolev space. Other results were obtained in similar frameworks ([Szabó et al., 2015](#); [Knapik et al., 2016](#); [Karvonen et al., 2020](#)).

This article presents three main contributions. First, we focus on the one-dimensional case of the framework proposed by [Stein \(1999, Section 6.7\)](#) to give an asymptotic normality result for maximum likelihood parameter estimation. Then, we leverage these convergence rates to analyze the expected integrated error, taking constant factors into account and showing that estimating the parameters yields the same error asymptotically as if the ground truth was known. Finally, we investigate the deterministic case by deriving the large sample limit of the likelihood criterion in a particular case. This suggests that bounding assumptions on some parameters can lead to different estimates.

The article is organized as follows. First, [Section 2.1](#) details the asymptotic framework and notations. Then, [Section 3](#) gives the main results. Finally, [Section 4](#) provides a few results on the deterministic case.

## 2 Gaussian process interpolation on the circle

### 2.1 Framework

Let  $f : [0, 1] \rightarrow \mathbb{R}$  be a continuous periodic function observed on a regular grid:

$$\{j/n, 0 \leq j \leq n-1\}.$$

Consider the periodic version of the stationary Matérn covariance function  $\{k_\theta, \theta \in (0, +\infty)^3\}$  proposed by [Stein \(1999, Section 6.7\)](#) and defined, for  $\theta = (\nu, \phi, \alpha) \in (0, +\infty)^3$ , by the uniformly absolutely convergent Fourier series with coefficients:

$$\underline{c}_j(\theta) = \frac{\phi}{(\alpha^2 + j^2)^{\nu+1/2}}, j \in \mathbb{Z}. \quad (1)$$

The parameter  $\alpha$  is not identifiable as different values yield equivalent probability measures. However,  $\nu$  and  $\phi$  are identifiable (see, e.g., [Stein, 1999, Chapter 4 and Section 6.7](#)). [Stein \(1999\)](#) advocates the *regularity* parameter  $\nu$  as the key quantity governing the asymptotics of the kriging error. The amplitude parameter  $\phi$  does not impact the kriging predictor but is known to be critical for uncertainty quantification (see, e.g., [Stein, 1993b](#)) and to be consistently estimated by maximum likelihood if  $\nu$  and  $\alpha$  are known (see, e.g., [Zhang, 2004](#), who use a different parametrization).

The usual task in kriging is to infer the function  $f$  from the data

$$Z = (f(0), \dots, f(1-1/n))^T.$$

---

<sup>1</sup>This framework is similar to that proposed by [Stein \(1993a\)](#) in a slightly different setting, where a different conjecture is made about the error on noisy training data.

## 2.2 Best linear prediction

The function  $f$  is usually predicted using the posterior mean function given by the kriging equations (Matheron, 1971). This predictor can be written simply in the framework presented above.

**Proposition 1.** *Let  $n \geq 1$  and  $f: [0, 1] \rightarrow \mathbb{R}$  a continuous periodic function. Suppose also that  $f$  is the pointwise absolute limit of its Fourier serie. Writing  $\widehat{f}_n$  for the posterior mean function given  $Z$  and the parameter  $\theta \in \Theta$ , we have:*

$$\widehat{f}_n(x) = \sum_{j \in \mathbb{Z}} \left( \frac{\sum_{j_1 \in j+n\mathbb{Z}} c_{j_1}(f)}{\sum_{j_1 \in j+n\mathbb{Z}} \underline{c}_{j_1}(\theta)} \right) \underline{c}_j(\theta) e^{2\pi i x j}, \quad x \in [0, 1], \quad (2)$$

where the  $c_j(f)$ s are the Fourier coefficients of  $f$ .

The expression (2) shows how the posterior mean function approximates  $f$ : it transforms the Fourier coefficients of  $k_\theta$  into those of  $f$  using the ratio of the discrete Fourier transforms. Finally, we also define the integrated squared error:

$$\text{ISE}_n(\mathbf{v}, \alpha; f) = \int_0^1 (f - \widehat{f}_n)^2. \quad (3)$$

Note that it does not depend on  $\phi$ .

## 2.3 Maximum likelihood estimation

Given the observations  $Z$  and  $\Theta \subset (0, +\infty)^3$ , the maximum likelihood estimate is defined by  $\widehat{\theta}_n = (\widehat{\mathbf{v}}_n, \widehat{\phi}_n, \widehat{\alpha}_n)$  minimizing (a linear transform of) the negative log-likelihood:

$$\mathbb{L}_n(\theta) = n^{-1} (\ln(\det(K_\theta)) + Z^\top K_\theta^{-1} Z), \quad \theta = (\mathbf{v}, \phi, \alpha) \in \Theta, \quad (4)$$

with ties broken arbitrarily and  $K_\theta$  the covariance matrix of  $Z$  under  $\theta$ .

The parameters  $\mathbf{v}$  and  $\alpha$  will be assumed to be bounded in this work, i.e., we take  $\Theta = N \times (0, +\infty) \times A$  with  $N$  and  $A$  bounded away from zero and infinity. Most of our results will be stated with  $A$  being a singleton, with  $\alpha$  enforced to a value (which will not necessarily be the ground truth). This type of assumption is more or less standard in the field, as it leads to simplifications (see, e.g., Ying, 1991; Loh, 2006; Chen et al., 2021). However, keeping  $\phi$  unbounded is key to our results and for discussing the deterministic case in Section 4.

## 2.4 A symmetrized version of the Hurwitz zeta function

As Stein (1999, Section 6.7) points out, a great simplification emerges when we restrict ourselves to the framework presented in Section 2.1. This will allow us to derive a sharp analysis, taking into account constant factors in the evaluation of the prediction accuracy with estimated parameters. Similarly to Stein (1999), the function

$$\gamma: (\alpha; x) \in (1, +\infty) \times (0, 1) \mapsto \sum_{j \in \mathbb{Z}} \frac{1}{|j+x|^\alpha},$$

will play a major role in deriving the limiting constants. It is  $C^\infty$  with respect to both arguments and related to the Hurwitz zeta function  $\zeta_H$  by:

$$\gamma(\alpha; x) = \zeta_H(\alpha; x) + \zeta_H(\alpha; 1-x), \quad x \in (0, 1). \quad (5)$$

Moreover, the function  $\gamma(\alpha; \cdot)$  is symmetric with respect to  $1/2$  for  $\alpha > 1$ .

Finally, the following will use the fact that  $\ln(\gamma(2\nu+1; \cdot))$ ,  $\gamma(2\nu_0+1; \cdot)/\gamma(2\nu+1; \cdot)$ , and  $\gamma^2(\nu_0+1; \cdot)/\gamma(2\nu+1; \cdot)$  are integrable for  $\nu > 0$ ,  $\nu > \nu_0 - 1/2$ , and  $\nu > \nu_0$ , respectively.

### 3 Main results

#### 3.1 Standing assumptions

Consider the framework presented in Section 2.1 and suppose that the function is sampled according to the (real-valued) centered Gaussian process:

$$\xi : x \in [0, 1] \mapsto \frac{1}{\sqrt{2}} \sum_{j \in \mathbb{Z}} \sqrt{\mathcal{L}_j(\theta_0)} (U_{1,|j|} + iU_{2,|j|}\text{sign}(j)) e^{2\pi i x j}, \quad (6)$$

with  $\theta_0 = (\nu_0, \phi_0, \alpha_0) \in (0, +\infty)^3$  and  $(U_{q,j})_{q \in \{1,2\}, j \geq 0}$  independent random variables such that  $U_{2,0} = 0$ ,  $U_{1,0} \sim \mathcal{N}(0, 2)$ , and  $U_{q,j} \sim \mathcal{N}(0, 1)$  for  $q \in \{1, 2\}$  and  $j \geq 1$ . It is easy to check that  $\xi \sim \text{GP}(0, k_{\theta_0})$ .

Let  $\hat{\theta}_n = (\hat{\nu}_n, \hat{\phi}_n, \hat{\alpha}_n)$  be a maximum likelihood estimate defined in Section 2.3 for some  $\Theta = N \times (0, +\infty) \times A$  with  $A \subset (0, +\infty)$  and  $N = [a_\nu, b_\nu] \ni \nu_0$ . The following sections give convergence rates in terms of parameter estimation and predictions.

#### 3.2 Convergence rate of maximum likelihood estimation

The following result states the consistency for the identifiable parameters and an upper bound of the corresponding rates.

**Theorem 1.** *Let  $0 < \beta < 1/4$  and  $A = [a_\alpha, b_\alpha]$  be bounded away from zero and infinity. The bounds  $\hat{\nu}_n - \nu_0 = o_P(n^{-\beta})$  and  $\hat{\phi}_n - \phi_0 = o_P(\ln(n)n^{-\beta})$  hold in probability.*

Informally, Theorem 1 is proven by showing that a linear transformation of

$$\inf_{\phi > 0} \mathbb{L}_n(\nu, \phi, \alpha)$$

converges to

$$\ln(\gamma(2\nu+1; \cdot)) + \ln\left(\int_0^1 \frac{\gamma(2\nu_0+1; \cdot)}{\gamma(2\nu+1; \cdot)}\right), \quad (7)$$

for  $\nu > \nu_0 - 1/2$ . Jensen inequality then shows that (7) is maximized by taking  $\nu = \nu_0$ .

Furthermore, similarly to Stein (1999, Section 6.7), let us define

$$\psi_\nu : x \in (0, 1) \mapsto \frac{\sum_{j \in \mathbb{Z}} |x+j|^{-2\nu-1} \ln|x+j|}{\sum_{j \in \mathbb{Z}} |x+j|^{-2\nu-1}}, \quad \text{for } \nu > 0,$$

which is square integrable on  $(0, 1)$  and  $e(\nu) = \mathbb{E}(\psi_\nu(U))$  and  $V(\nu) = \text{Var}(\psi_\nu(U))$ , with  $U \sim \mathcal{U}(0, 1)$ . The following result proves the conjecture made by Stein (1999, Section 6.7) when  $\hat{\nu}_n$  is bounded and  $\alpha_0$  is known.

**Theorem 2.** *Suppose that  $A = \{\alpha_0\}$ . Then*

$$\sqrt{2n} \left( \frac{\hat{\phi}_n - \phi_0}{2\phi_0} - (\ln(n) + e(\nu_0)) (\hat{\nu}_n - \nu_0) \right) \rightsquigarrow \mathcal{N}(0, I_2). \quad (8)$$

### 3.3 Convergence rate of the integrated prediction error

This section states our results about the expectation of (3) with estimated parameters. To avoid technical difficulties, we suppose that  $\nu_0 > 1/2$  in this section. We begin with the case of fixed parameters.

For  $\nu, \nu_0 > 0$ , define

$$\vartheta_{\nu, \nu_0} : x \in (0, 1) \mapsto \frac{\gamma(4\nu + 2; x) \gamma(2\nu_0 + 1; x)}{\gamma^2(2\nu + 1; x)} + \gamma(2\nu_0 + 1; x) - 2 \frac{\gamma(2\nu + 2\nu_0 + 2; x)}{\gamma(2\nu + 1; x)}.$$

The function  $\vartheta_{\nu, \nu_0}$  is  $C^1$  on  $(0, 1)$ , symmetric with respect to  $1/2$ , and a Taylor expansion around  $x = 0$  shows that it is integrable if  $\nu > (\nu_0 - 1)/2$ .

The following result states the asymptotics of the prediction error with fixed parameters.

**Theorem 3.** *Let  $\theta = (\nu, \phi, \alpha) \in N \times \mathbb{R} \times A$ . Then,*

$$\mathbb{E}(\text{ISE}_n(\nu, \alpha; \xi)) \lesssim \frac{1}{n^{4\nu+2}}, \text{ for } \nu < (\nu_0 - 1)/2 \quad (9)$$

$$\mathbb{E}(\text{ISE}_n(\nu, \alpha; \xi)) \lesssim \frac{\ln(n)}{n^{2\nu_0}}, \text{ for } \nu = (\nu_0 - 1)/2 \quad (10)$$

and

$$\mathbb{E}(\text{ISE}_n(\nu, \alpha; \xi)) \sim \frac{\phi_0 \int_0^1 \vartheta_{\nu, \nu_0}}{n^{2\nu_0}}, \text{ otherwise.} \quad (11)$$

The symbol  $\lesssim$  denotes an inequality up to a universal constant.

This result shows that half of the smoothness is sufficient for optimal convergence rates. However, the constant  $\int_0^1 \vartheta_{\nu, \nu_0}$  is minimized by taking  $\nu = \nu_0$ . This is in line with the result of Stein (1999, Theorem 3) obtained in a different framework.

Then, our last result gives the asymptotic behavior of the prediction error with estimated  $\nu$  and  $\phi$  and fixed—but not necessarily known— $\alpha$ .

**Theorem 4.** *Let  $A = \{\alpha\}$  with  $0 < \alpha < +\infty$ . Then,*

$$\mathbb{E}(\text{ISE}_n(\hat{\nu}_n, \alpha; \xi)) \sim \frac{\phi_0 \int_0^1 \vartheta_{\nu_0, \nu_0}}{n^{2\nu_0}}. \quad (12)$$

This last result shows that estimating the parameters is optimal both in terms of the rate and the constant.

## 4 The deterministic case

This section studies the case of an unknown fixed deterministic function from a continuous Sobolev space. Let  $\beta > 0$  and define the Sobolev space

$$H_{[0,1]}^{\beta+1/2} = \left\{ g \in L_{[0,1]}^2, \|g\|_{H_{[0,1]}^{\beta+1/2}}^2 = \sum_{j \in \mathbb{Z}} (1+j^2)^{\beta+1/2} |c_j(f)|^2 < +\infty \right\} \quad (13)$$

of (continuous) periodic functions.

Furthermore, let  $f \in H_{[0,1]}^{\beta+1/2}$  for some  $\beta > 0$  and define the smoothness

$$v_0(f) = \inf \left\{ \beta > 0, f \notin H_{[0,1]}^{\beta+1/2} \right\}$$

of  $f$  as Wang and Jing (2022) and Karvonen (2022).

Suppose that  $\hat{\theta}_n = (\hat{v}_n, \hat{\phi}_n, \hat{\alpha}_n)$  is estimated by maximizing the likelihood as in Section 2.3 for  $\Theta = N \times F \times A$  with  $N$  and  $A$  bounded away from zero and infinity. This section discusses the behavior of  $\hat{v}_n$  under three assumptions on  $F$ : 1) a singleton; 2) a range bounded away from zero and infinity; and 3) the whole  $(0, +\infty)$ . For the last case, the definition

$$\mathbb{M}_n^f: (v, \alpha) \in N \times A \mapsto \inf_{\phi > 0} \mathbb{L}_n(v, \phi, \alpha) + (2v_0(f) + 1) \ln(n) - 1,$$

will be used.

On “nice” bounded regions of  $\mathbb{R}^d$ , Karvonen (2022) shows that  $\liminf \hat{v}_n \geq v_0(f)$  if  $\alpha$  and  $\phi$  are fixed<sup>2</sup>. The following result shows that it holds on the circle no matter the assumption on  $F$ .

**Proposition 2.** *If  $N$  and  $A$  are bounded away from zero and infinity, then*

$$\liminf \hat{v}_n \geq v_0(f)$$

*holds for the three previous assumptions on  $F$ .*

Regarding the precise behavior of  $\hat{v}_n$  above  $v_0(f)$ , Karvonen (2022) conjectures that  $\hat{v}_n$  converges to  $v_0(f) + 1/2$  if  $\phi$  and  $\alpha$  are fixed but deems that a joint estimation may give a different behavior. The rest of this section is devoted to supporting this idea.

Write  $\approx$  for a two-way inequality up to universal constants and use the more stringent assumption that

$$|c_j(f)| \approx |j|^{-v_0(f)-1}$$

for simplicity. The following result is a minor adaptation of the reasoning used by Chen et al. (2021) and Karvonen (2022), showing that the conjecture is verified more generally if  $\phi$  is bounded.

**Proposition 3.** *Suppose that  $|c_j(f)| \approx |j|^{-v_0(f)-1}$  and that  $N$ ,  $A$ , and  $F$  are bounded away from zero and infinity. Then, the convergence  $\hat{v}_n \rightarrow v_0(f) + 1/2$  holds.*

However, our last results suggest that it does not hold when  $F = (0, +\infty)$ .

**Proposition 4.** *Suppose that  $c_j(f) = |j|^{-v_0(f)-1}$ , for  $j \neq 0$ . Then, for  $v > v_0(f)$  and  $\alpha > 0$ , we have*

$$\mathbb{M}_n^f(v, \alpha) \rightarrow \mathbb{M}_\infty^f(v) = \int_0^1 \ln(\gamma(2v+1; \cdot)) + \ln \left( \int_0^1 \frac{\gamma^2(v_0(f)+1; \cdot)}{\gamma(2v+1; \cdot)} \right).$$

It is possible to obtain a uniform version of the previous convergence. However, it is omitted for brevity and because we could not identify the minimizer(s) of the limit analytically. Figure 1 shows a numerical approximation of  $\mathbb{M}_\infty^f$ .

<sup>2</sup>Note that Karvonen (2022) does not suppose that  $v$  is bounded.




Figure 1: The function  $M_\infty^f$ , for  $v_0(f) = 1$ . A numerical approximation of the minimizer is about 1.359. Note that  $M_\infty^f$  is approximated numerically using finite sums for  $\gamma$  and discretizations for the integrals.



## A Proofs

### A.1 Additional notations

The symbol  $\lesssim$  denotes an inequality up to a universal constant. For compactness, the symbol  $\approx$  is used when the two-way inequality  $\lesssim$  holds.

Write  $K_\theta = \phi R_{v,\alpha}$  and  $\underline{c}_j(\theta) = \phi c_j(v, \alpha)$ , for  $\theta = (v, \phi, \alpha) \in (0, +\infty)^3$  and  $j \in \mathbb{Z}$ . All results suppose that  $\Theta = N \times (0, +\infty) \times A$  with  $N = [a_v, b_v]$ ,  $A = [a_\alpha, b_\alpha]$ ,  $0 < a_v \leq b_v < +\infty$ ,  $0 < a_\alpha \leq b_\alpha < +\infty$ , and  $F = (0, +\infty)$  unless explicitly stated otherwise. Without loss of generality, suppose that  $a_v < v_0 - 1/2$  and define  $N_\varepsilon = [v_0 - 1/2 + \varepsilon, b_v] \subset N$  for  $\varepsilon > 0$ .

Furthermore, for  $\theta_0 = (v_0, \phi_0, \alpha_0)$ , define  $\mathbb{M}_n$  to be the stochastic process:

$$\mathbb{M}_n: (v, \alpha) \in N \times A \mapsto \inf_{\phi > 0} \mathbb{L}_n(v, \phi, \alpha) + 2v_0 \ln(n) - 1.$$

Its expression is given almost surely by the following Lemma.

**Lemma 1.** (see, e.g., [Santner et al., 2003](#), Section 3.3.2) *Let  $v, \alpha > 0$ . If  $Z \neq 0$ , then*

$$\inf_{\phi > 0} \mathbb{L}_n(v, \phi, \alpha) = 1 + n^{-1} \ln(\det(R_{v,\alpha})) + \ln\left(\frac{Z^\top R_{v,\alpha}^{-1} Z}{n}\right), \quad (14)$$

where the supremum is reached by  $\hat{\phi}_n = Z^\top R_{v,\alpha}^{-1} Z / n$ .

Note that  $\mathbb{M}_n$  has therefore almost surely continuous sample paths. The proofs mostly consist in studying  $\mathbb{M}_n$ , which is often called the profiled likelihood. For a compact  $A \subset (0, +\infty)$ , define now  $\mathbb{U}_n: v \in N \mapsto \inf_{\alpha \in A} \mathbb{M}_n(v, \alpha)$ . The object  $\mathbb{U}_n$  is still a collection of measurable maps and is also almost surely continuous since  $A$  is compact.

Also, define

$$g_v = \ln(\gamma(2v + 1; \cdot)),$$

for  $v > 0$  and

$$h_{v,v_0} = \frac{\gamma(2v_0 + 1; \cdot)}{\gamma(2v + 1; \cdot)},$$

for  $v > v_0 - 1/2$ . These functions are  $C^\infty$  and integrable.

Finally, suppose  $n \geq 3$  and use the notation  $l = \lfloor (n-1)/2 \rfloor$  throughout the following.

### A.2 Circulant matrices and useful facts

Let us remind how the framework of Section 2.1 yields simplifications that can be traced back at least to [Craven and Wahba \(1979\)](#).

First let  $W$  be the  $n \times n$  matrix with entries  $W_{j,m} = \sqrt{n}^{-1} e^{2\pi i j m / n}$ , for  $0 \leq j, m \leq n-1$ . For every  $\theta = (v, \phi, \alpha) \in (0, +\infty)^3$ , the matrix  $R_{v,\alpha}$  is circulant and thus  $R_{v,\alpha} = W \Delta_{v,\alpha} W^*$  (see, e.g., [Brockwell and Davis, 1987](#), p. 130) with

$$\Delta_{v,\alpha} = \text{diag}(\lambda_{0,n}, \dots, \lambda_{n-1,n})$$

and

$$\lambda_{m,n} = \sum_{j=0}^{n-1} e^{-2\pi i j m / n} k_\theta(j/n) = n \sum_{j \in \mathbb{Z}} c_{m+nj}(\theta), \quad 0 \leq m \leq n-1. \quad (15)$$

These coefficients verify

$$\lambda_{m,n} = \lambda_{n-m,n}, \text{ for } 0 \leq m \leq n-1. \quad (16)$$

Furthermore, the matrices  $R_{\mathbf{v},\alpha}$  are symmetric so one also has  $R_{\mathbf{v},\alpha} = P\Delta P^\top$  for a unitary matrix  $P$ , obtained, e.g., by taking the real and imaginary parts of  $W$ . For  $\theta_0 = (\mathbf{v}_0, \phi_0, \alpha_0)$ , write

$$P^\top Z = \sqrt{\phi_0} (\lambda_{0,n}^0 U_{0,n}, \dots, \lambda_{n-1,n}^0 U_{n-1,n}),$$

with  $\lambda_{0,n}^0, \dots, \lambda_{n-1,n}^0$  the eigenvalues of  $R_{\mathbf{v}_0, \alpha_0}$  and  $U_{0,n}, \dots, U_{n-1,n}$  drawn independently from a standard Gaussian. We have

$$Z^\top R_{\mathbf{v},\alpha}^{-1} Z = \phi_0 \sum_{m=0}^{n-1} \frac{U_{m,n}^2 \lambda_{m,n}^0}{\lambda_{m,n}}. \quad (17)$$

### A.3 Proofs of Section 2.2

*Proof of Proposition 1.* The kriging equations yield  $\hat{f}_n(x) = k_{\theta,x}^\top K_\theta^{-1} Z$ , with  $k_{\theta,x} = (k_\theta(m/n - x))_{0 \leq m \leq n-1}$ , for  $x \in [0, 1]$ . The simplifications described in Section A.2 give:

$$W^* Z = \sqrt{n} \left( \sum_{j \in m+n\mathbb{Z}} c_j(f) \right)_{0 \leq m \leq n-1},$$

and

$$W^* k_{\theta,x} = \sqrt{n} \left( \sum_{j \in m+n\mathbb{Z}} \underline{c}_j(\theta) e^{-2\pi i x j} \right)_{0 \leq m \leq n-1}.$$

The result follows from elementary manipulations.  $\square$

### A.4 Proof of Theorem 1

**Lemma 2.** *The quantities  $\lambda_{0,n}/n$  and  $\lambda_{m,n} m^{2v+1}/n$  are bounded away from zero and infinity uniformly in  $\mathbf{v} \in N$ ,  $\alpha \in A$ ,  $n$  and  $1 \leq m \leq \lfloor n/2 \rfloor$ .*

*Proof.* Let  $0 \leq m \leq \lfloor n/2 \rfloor$ , we have using (15)

$$c_m(\mathbf{v}, \alpha) \leq \lambda_{m,n}/n \leq 2c_m(\mathbf{v}, \alpha) + 2 \sum_{j=1}^{+\infty} c_{m+nj}(\mathbf{v}, \alpha).$$

Moreover

$$\sum_{j=1}^{+\infty} c_{m+nj}(\mathbf{v}, \alpha) / c_m(\mathbf{v}, \alpha) \leq \sum_{j=1}^{+\infty} (b_\alpha^2 + 1/4)^{v+1/2} / j^{2v+1} \leq C_1^{b_\alpha, a_v, b_v} \quad (18)$$

where  $C_1^{b_\alpha, a_v, b_v} = \max(1, (b_\alpha^2 + 1/4)^{b_v+1/2}) \zeta(2a_v + 1)$ . The result follows from elementary operations.  $\square$

**Lemma 3.** *Let  $\mathbf{v} \in N$ ,  $\alpha \in A$ , and  $1 \leq m \leq l$ , we have*

$$c_{m+nj}(\mathbf{v}, \alpha) = \frac{(1 + u_{n,m,j}(\mathbf{v}, \alpha))}{|jn + m|^{2v+1}}, \quad (19)$$

with  $-1 < v_m \leq u_{n,m,j}(\mathbf{v}, \alpha) \leq 0$  and  $v_m = \mathcal{O}(m^{-2})$ .

*Proof.* Using (15), we have

$$\begin{aligned} c_{m+nj}(\mathbf{v}, \alpha) &= \frac{1}{(\alpha^2 + (jn+m)^2)^{\nu+1/2}} \\ &= \frac{(1 + u_{n,m,j}(\mathbf{v}, \alpha))}{|jn+m|^{2\nu+1}}, \end{aligned}$$

with  $u_{n,m,j}(\mathbf{v}, \alpha) = (1 + (\alpha/(jn+m))^2)^{-\nu-1/2} - 1$ . Elementary operations show that

$$0 \geq u_{n,m,j}(\theta_1) \geq \left( \left( \frac{b\alpha}{m} \right)^2 + 1 \right)^{-b\nu-1/2} - 1,$$

which gives the desired result thanks to the Taylor inequality.  $\square$

**Lemma 4.** *Uniformly in  $\mathbf{v} \in N$  and  $\alpha \in A$ , we have*

$$\ln(\det(R_{\mathbf{v},\alpha})) = -2\nu n \ln(n) + n \int_0^1 g_{\mathbf{v}}(x) dx + \mathcal{O}(\ln(n)).$$

*Proof.* Using (15) and Lemma 3, we have

$$\begin{aligned} \lambda_{m,n}/n &= \sum_{j \in \mathbb{Z}} c_{m+nj}(\mathbf{v}, \alpha) \\ &= \sum_{j \in \mathbb{Z}} \frac{(1 + u_{n,m,j}(\mathbf{v}, \alpha))}{|jn+m|^{2\nu+1}}, \end{aligned}$$

and therefore

$$\sum_{m=1}^l \ln(\lambda_{m,n}/n) = -(2\nu+1)l \ln(n) + a_n + \sum_{m=1}^l g_{\mathbf{v}}(m/n),$$

with

$$|a_n| \leq \left| \sum_{m=1}^l \ln(1 + v_m) \right|$$

which is thus  $\mathcal{O}(1)$  uniformly in  $\mathbf{v} \in N$  and  $\alpha \in A$ .

One has  $g_{\mathbf{v}}(x) = g_{\mathbf{v}}(1-x)$  for  $x \in (0,1)$  and it is easy to see that  $g_{\mathbf{v}}$  is non-increasing on  $(0, 1/2]$ , so

$$\int_{1/n}^{(l+1)/n} g_{\mathbf{v}}(x) dx \leq \frac{1}{n} \sum_{m=1}^l g_{\mathbf{v}}(m/n) \leq \int_0^{l/n} g_{\mathbf{v}}(x) dx.$$

First we have  $\left| \int_{l/n}^{1/2} g_{\mathbf{v}} \right| \leq |g_{\mathbf{v}}(1/4)|/(2n) = \mathcal{O}(n^{-1})$  uniformly in  $\mathbf{v} \in N$  thanks to a compactness argument and similarly for  $\left| \int_{(l+1)/n}^{1/2} g_{\mathbf{v}} \right|$ .

Moreover observe that

$$\gamma_{2\nu+1}(x) - 1/x^{2\nu+1} = \sum_{j \in \mathbb{Z}, j \neq 0} \frac{1}{|j+x|^{2\nu+1}}$$

is uniformly bounded in  $v \in N$  and thus that the equivalence

$$\mathcal{Y}_{2v+1}(x) \sim_{x \rightarrow 0} 1/x^{2v+1} \quad (20)$$

is uniform and so is

$$g_v(x) \sim_{x \rightarrow 0} -(2v+1) \ln(x).$$

Therefore, we have

$$\int_0^{1/n} g_v(x) dx = \mathcal{O}(\ln(n)/n),$$

uniformly in  $v \in N$  and so

$$\sum_{m=1}^l g_v(m/n) = n \int_0^{1/2} g_v(x) dx + \mathcal{O}(\ln(n)).$$

Moreover, Lemma 2 shows that  $\ln(\lambda_{0,n}/n) = \mathcal{O}(1)$  and  $\ln(\lambda_{n/2,n}/n) = \mathcal{O}(\ln(n))$  uniformly for  $n$  even. One can then conclude using (16).  $\square$

**Lemma 5.** *The function  $h_{v;v_0}$  is non-decreasing (resp. non-increasing) on  $(0, 1/2]$  when  $v \geq v_0$  (resp.  $v \leq v_0$ ).*

*Proof.* Suppose that  $v \geq v_0$ . Use (5) along with the fact that the Hurwitz Zeta function verifies (Postnikov, 1988)

$$\frac{\partial \zeta_H}{\partial x}(\alpha; x) = -\alpha \zeta_H(\alpha+1; x), \quad x > 0, \alpha > 1 \quad (21)$$

and has the representation

$$\zeta_H(\alpha; x) = \frac{1}{\Gamma(\alpha)} \int_0^{+\infty} \frac{t^{\alpha-1} e^{-tx}}{1-e^{-t}} dt, \quad x > 0, \alpha > 1,$$

where  $\Gamma$  is the classical Gamma function. So we have

$$\mathcal{Y}_{2v+1}(x) = \frac{1}{\Gamma(2v+1)} \int_0^{+\infty} \frac{t^{2v} (e^{-tx} + e^{-t(1-x)})}{1-e^{-t}} dt, \quad x \in (0, 1),$$

and

$$\frac{d\mathcal{Y}_{2v+1}}{dx}(x) = \frac{1}{\Gamma(2v+1)} \int_0^{+\infty} \frac{t^{2v+1} (e^{-t(1-x)} - e^{-tx})}{1-e^{-t}} dt, \quad x \in (0, 1).$$

Now let  $x \in [1/2, 1)$ , the derivative of  $h_{v;v_0}$  at  $x$  has the sign of

$$\begin{aligned} & \mathcal{Y}_{2v+1}(x) \frac{d\mathcal{Y}_{2v_0+1}}{dx}(x) - \mathcal{Y}_{2v_0+1}(x) \frac{d\mathcal{Y}_{2v+1}}{dx}(x) \\ &= \frac{1}{\Gamma(2v+1)\Gamma(2v_0+1)} \int_0^{+\infty} \int_0^{+\infty} \frac{t^{2v} s^{2v_0} (\eta(s, t; x) - \eta(t, s; x))}{\kappa(s, t)} dt ds \end{aligned}$$

with  $\eta(s, t; x) = s(e^{-tx} + e^{-t(1-x)})(e^{-s(1-x)} - e^{-sx})$  and  $\kappa(s, t) = (1-e^{-t})(1-e^{-s}) = \kappa(t, s)$  thanks to the Fubini-Lebesgue theorem. Then, one can split the integral to have:

$$\frac{1}{\Gamma(2v+1)\Gamma(2v_0+1)} \left( \int_0^{+\infty} \int_t^{+\infty} \frac{t^{2v} s^{2v_0} (\eta(s, t; x) - \eta(t, s; x))}{\kappa(s, t)} dt ds \right)$$

$$\begin{aligned}
& + \int_0^{+\infty} \int_t^{+\infty} \frac{s^{2\nu} t^{2\nu_0} (\eta(t, s; x) - \eta(s, t; x))}{\kappa(t, s)} dt ds \\
& = \frac{1}{\Gamma(2\nu + 1) \Gamma(2\nu_0 + 1)} \int_0^{+\infty} \int_t^{+\infty} \frac{(t^{2\nu} s^{2\nu_0} - s^{2\nu} t^{2\nu_0}) (\eta(s, t; x) - \eta(t, s; x))}{\kappa(s, t)} dt ds \leq 0
\end{aligned}$$

since  $t^{2\nu} s^{2\nu_0} \leq s^{2\nu} t^{2\nu_0}$  when  $s \geq t$ ,  $\kappa(s, t) \geq 0$  and  $\eta(s, t; x) \geq \eta(t, s; x)$  when  $s \geq t$  and  $x \geq 1/2$ .

So we proved that  $h_{\nu, \nu_0}$  is non-increasing on  $[1/2, 1)$  and the first claim is due to the symmetry with respect to  $1/2$ . Observe that  $h_{\nu, \nu_0} = 1/h_{\nu_0, \nu}$  for the second claim.  $\square$

**Lemma 6.** *Let  $\varepsilon > 0$ , we have*

$$\frac{1}{n} \sum_{m=1}^{n-1} h_{\nu, \nu_0}(m/n) = \int_0^1 h_{\nu, \nu_0}(x) dx + \mathcal{O}\left(\frac{1}{n^{\min(1, 2\varepsilon)}}\right),$$

uniformly in  $\nu \in N_\varepsilon$ .

*Proof.* Using Lemma 5 and the symmetry w.r.t.  $1/2$ , we have again the classical

$$\int_{1/n}^{(l+1)/n} h_{\nu, \nu_0}(x) dx \leq \frac{1}{n} \sum_{m=1}^l h_{\nu, \nu_0}(m/n) \leq \int_0^{l/n} h_{\nu, \nu_0}(x) dx, \quad (22)$$

when  $\nu \geq \nu_0$  and the reversed inequality when  $\nu \leq \nu_0$ .

The behaviour near  $1/2$  is elucidated as for Lemma 4 with a compacity argument to bound the values of  $h_{\nu, \nu_0}$  uniformly at  $1/4$  and  $1/2$ , thus giving a uniform  $\mathcal{O}(n^{-1})$ .

Moreover, it is straightforward to show that

$$h_{\nu, \nu_0}(x) \sim_{x \rightarrow 0} x^{2(\nu - \nu_0)}, \quad (23)$$

uniformly in  $\nu \in N$  and then

$$\int_0^{1/n} h_{\nu, \nu_0}(x) dx = \mathcal{O}(n^{-2\varepsilon}), \quad (24)$$

uniformly in  $\nu \in N_\varepsilon$ . Therefore, we have

$$\sum_{m=1}^l h_{\nu, \nu_0}(m/n) = n \int_0^{1/2} h_{\nu, \nu_0}(x) dx + \mathcal{O}(1) + \mathcal{O}(n^{1-2\varepsilon}),$$

uniformly and using the symmetry of  $h_{\nu, \nu_0}$  gives the desired result.  $\square$

**Lemma 7.** *Let  $1 \leq m \leq l$ , we have*

$$\frac{\lambda_{m,n}^0}{n^{2(\nu - \nu_0)} \lambda_{m,n}} = (1 + \mathcal{O}(m^{-2})) h_{\nu, \nu_0}(m/n)$$

uniformly in  $\nu \in N$  and  $\alpha \in A$ .

*Proof.* Lemma 3 gives

$$\frac{\lambda_{m,n}^0}{n^{2(\nu - \nu_0)} \lambda_{m,n}} = \frac{\sum_{j \in \mathbb{Z}} \frac{(1 + u_{n,m,j}(\nu_0, \alpha_0))}{|j + m/n|^{2\nu_0 + 1}}}{\sum_{j \in \mathbb{Z}} \frac{(1 + u_{n,m,j}(\nu, \alpha))}{|j + m/n|^{2\nu + 1}}},$$

with

$$\sum_{j \in \mathbb{Z}} \frac{(1 + u_{n,m,j}(\nu, \alpha))}{|j + m/n|^{2\nu + 1}} = \gamma_{2\nu+1}(m/n) (1 + \mathcal{O}(m^{-2})),$$

uniformly. The desired result follows then from elementary manipulations.  $\square$

**Lemma 8.** *Let  $0 < \varepsilon < 1/2$ , we have*

$$\sup_{(\nu, \alpha) \in N_\varepsilon \times A} \frac{1}{n} \left| \sum_{m=1}^{n-1} \left( \frac{\lambda_{m,n}^0}{n^{2(\nu-\nu_0)} \lambda_{m,n}} - h_{\nu; \nu_0} \left( \frac{m}{n} \right) \right) \right| = \mathcal{O} \left( \frac{1}{n^{\min(3/4, 5\varepsilon/4)}} \right).$$

*Proof.* Let  $1 \leq m \leq l$  and  $\beta \in (0, 1)$ . Using Lemma 7 and the correct sign for the big- $\mathcal{O}$  term we have

$$\begin{aligned} \frac{1}{n} \left| \sum_{m=1}^l \left( \frac{\lambda_{m,n}^0}{n^{2(\nu-\nu_0)} \lambda_{m,n}} - h_{\nu; \nu_0} \left( \frac{m}{n} \right) \right) \right| &\leq \frac{1}{n} \sum_{m=1}^l \mathcal{O} \left( m^{-2} \right) h_{\nu; \nu_0} \left( \frac{m}{n} \right) \\ &= \frac{1}{n} \sum_{m=1}^{\lfloor n^\beta \rfloor} \mathcal{O} \left( m^{-2} \right) h_{\nu; \nu_0} \left( \frac{m}{n} \right) + \frac{1}{n} \sum_{m=\lfloor n^\beta \rfloor + 1}^l \mathcal{O} \left( m^{-2} \right) h_{\nu; \nu_0} \left( \frac{m}{n} \right), \end{aligned}$$

with

$$\frac{1}{n} \sum_{m=\lfloor n^\beta \rfloor + 1}^l \mathcal{O} \left( m^{-2} \right) h_{\nu; \nu_0} \left( \frac{m}{n} \right) = \mathcal{O} \left( \frac{1}{n^{2\beta}} \right) \frac{1}{n} \sum_{m=1}^l h_{\nu; \nu_0} \left( \frac{m}{n} \right) = \mathcal{O} \left( \frac{1}{n^{2\beta}} \right)$$

uniformly in  $\nu \in N_\varepsilon$  and  $\alpha \in A$  thanks to (22) and by observing that the mapping  $\nu \in N_\varepsilon \mapsto \int_0^{1/2} h_{\nu; \nu_0}$  is continuous. Moreover we have

$$\sum_{m=1}^{\lfloor n^\beta \rfloor} \mathcal{O} \left( m^{-2} \right) h_{\nu; \nu_0} \left( \frac{m}{n} \right) \leq \mathcal{O}(1) n \int_0^{\frac{\lfloor n^\beta \rfloor + 1}{n}} h_{\nu; \nu_0}(x) dx = \mathcal{O} \left( \frac{1}{n^{2\varepsilon(1-\beta)}} \right)$$

uniformly thanks to (22) and (24). Using (16), the symmetry of  $h_{\nu; \nu_0}$  and taking  $\beta = 3/8$  then gives the desired result.  $\square$

**Lemma 9.** *The family  $\mathcal{F} = \{h_{\nu; \nu_0}, \nu \in N\}$  satisfies the uniformly integrable entropy condition (Ziegler, 1997).*

*Proof.* We have  $h_{\nu; \nu_0} = \gamma_{\nu_0+1} / \gamma_{\nu+1}$ , with

$$\gamma_{2\nu+1}(x) = \sum_{j=0}^{+\infty} \frac{1}{(j+1+x)^{2\nu+1}} + \sum_{j=0}^{+\infty} \frac{1}{(j+2-x)^{2\nu+1}} + \frac{1}{x^{2\nu+1}} + \frac{1}{(1-x)^{2\nu+1}}, \quad 0 < x < 1.$$

The four families of functions involved in the last expression of  $\gamma_{2\nu+1}$  are monotonous with respect to  $\nu \in N$  so they are VC-subgraph classes of functions (Van Der Vaart and Wellner, 1996, Section 2.6.2). Indeed let  $(x_1, y_1), (x_2, y_2) \in (0, 1) \times \mathbb{R}$ , there cannot be  $f, g \in \mathcal{F}$  such that  $f(x_1) < y_1, f(x_2) \geq y_2, g(x_1) \geq y_1$  and  $g(x_2) < y_2$  since we have either  $g \leq f$  or  $f \leq g$ .

The class  $\mathcal{F}$  is therefore a VC-subgraph class of functions thanks to the Lemma 2.6.18 from Van Der Vaart and Wellner (1996) by observing that the statement (viii) remains valid for a monotonous function  $\phi$  defined only on a subset of  $\mathbb{R}$  containing the ranges of the elements of  $\mathcal{F}$ . Therefore  $\mathcal{F}$  satisfies the uniformly integrable entropy condition thanks to the Lemma 2.4 from Ziegler (1997), observing that it relies on a definition of the VC dimension that is equivalent (Van Der Vaart and Wellner, 1996, Exercice 2.6.11).  $\square$

**Lemma 10.** *We have*

$$\frac{1}{n} \sum_{m=0}^{n-1} (h_{v_1;v_0}(m/n) - h_{v_2;v_0}(m/n))^2 \rightarrow \int_0^1 (h_{v_1;v_0} - h_{v_2;v_0})^2,$$

uniformly in  $v_1, v_2 \in [v_0 - 1/4, b_v]$ .

*Proof.* Let  $\varepsilon > 0$ , we have

$$\int_0^\alpha (h_{v_1;v_0} - h_{v_2;v_0})^2 \leq 4 \int_0^\alpha F_\varepsilon^2 \leq \varepsilon/5$$

and

$$\frac{1}{n} \sum_{m=1}^{\lfloor \alpha n \rfloor} (h_{v_1;v_0}(m/n) - h_{v_2;v_0}(m/n))^2 \leq \frac{4}{n} \sum_{m=1}^{\lfloor \alpha n \rfloor} F_\varepsilon^2(m/n) \leq \varepsilon/5$$

and similarly for  $\frac{1}{n} \sum_{m=\lfloor (1-\alpha)n \rfloor}^{n-1} (h_{v_1;v_0}(m/n) - h_{v_2;v_0}(m/n))^2$  for some  $\alpha > 0$  and  $n$  that is large enough. Now a compactness argument shows that the derivative of  $(h_{v_1;v_0} - h_{v_2;v_0})^2$  is uniformly bounded on  $[\alpha, 1 - \alpha]$  so an adaptation of the traditional technique for error bounds on Riemann sums give

$$\left| \frac{1}{n} \sum_{m=\lfloor \alpha n \rfloor+1}^{\lfloor (1-\alpha)n \rfloor-1} (h_{v_1;v_0}(m/n) - h_{v_2;v_0}(m/n))^2 - \int_\alpha^{1-\alpha} (h_{v_1;v_0} - h_{v_2;v_0})^2 \right| \leq \varepsilon/5,$$

uniformly in  $v_1, v_2 \in [v_0 - 1/4, b_v]$  for  $n$  large enough. □

For  $n \geq 3$  and  $0 \leq m \leq n-1$ , define  $B_{m,n} = U_{m,n}^2 - 1$ .

**Lemma 11.** *Let  $3/8 < \varepsilon < 1/2$ , we have*

$$\frac{1}{\sqrt{n}} \sum_{m=1}^{n-1} B_{m,n} h_{\cdot;v_0} \left( \frac{m}{n} \right) \rightsquigarrow \text{GP} \left( 0, (v_1, v_2) \mapsto \int_0^1 h_{v_1;v_0} h_{v_2;v_0} \right),$$

in  $L^\infty(N_\varepsilon)$ .

*Proof.* First Lemma 9 shows that the family of functions  $\{h_{v;v_0}, v \in N_\varepsilon\}$  verifies the uniformly integrable entropy condition. Moreover, (23) gives the  $L_{(0,1)}^4$  integrable envelope

$$\sup_{v \in N_\varepsilon} h_{v;v_0}(x) \leq C \min(x, 1-x)^{-1+2\varepsilon} = F_\varepsilon(x), \quad 0 < x < 1,$$

for some  $C > 0$ .

The standard approach for monotonous Riemann sums applies by splitting on  $(0, 1/2]$  and  $[1/2, 1)$  and we have  $\sum_{m=1}^{n-1} F_\varepsilon^4(m/n) = \mathcal{O}(n)$  and therefore

$$\sum_{m=1}^{n-1} \mathbb{P} \left( \frac{F_\varepsilon(m/n) |B_{m,n}|}{\sqrt{n}} > \theta_n \right) \leq \frac{\mathbb{E}(B_{0,1}^4)}{n^2 \theta_n^4} \sum_{m=1}^{n-1} F_\varepsilon^4(m/n) = \mathcal{O} \left( \frac{1}{n \theta_n^4} \right),$$

which converges to zero with  $\theta_n = n^{-1/5}$ .

Moreover let  $d = \|\cdot - \cdot\|_{L^2_{(0,1)}}$ , one has  $\|F\|_{L^2_{(0,1)}} < +\infty$  so  $(\mathcal{F}, d)$  is totally bounded since it is covered by the uniformly integrable entropy condition. Let  $\beta, \rho > 0$ , we have

$$\begin{aligned} a_n(\beta, \rho) &= \sup_{d(f, g) \leq \beta} \mathbb{E} \left( \left( \sum_{m=1}^{n-1} (f(m/n) - g(m/n)) \frac{B_{m,n}}{\sqrt{n}} \mathbb{1}_{F_\varepsilon(m/n) |B_{m,n}| \leq \sqrt{n}\theta_n} \right)^2 \right) \\ &\leq \sup_{d(f, g) \leq \beta} \mathbb{E} \left( \left( \sum_{m=1}^{n-1} (f(m/n) - g(m/n)) \frac{B_{m,n}}{\sqrt{n}} \right)^2 \right) \\ &= \frac{\text{Var}(B_{0,1})}{n} \sup_{d(f, g) \leq \beta} \sum_{m=1}^{n-1} (f(m/n) - g(m/n))^2 \rightarrow \text{Var}(B_{0,1}) \sup_{d(f, g) \leq \beta} d^2(f, g). \end{aligned}$$

thanks to Lemma 10. Finally

$$\sup_{n \geq 1} \sum_{m=0}^{n-1} \mathbb{E} \left( F_\varepsilon^2(m/n) \frac{B_{m,n}^2}{n} \mathbb{1}_{F_\varepsilon(m/n) |B_{m,n}| \leq \sqrt{n}\theta_n} \right) \leq \sup_{n \geq 1} \frac{\text{Var}(B_{0,1})}{n} \sum_{m=0}^{n-1} F_\varepsilon^2(m/n),$$

is finite.

Now the finite dimensional laws convergence is stated using a classical Lyapunov condition observing that the  $B_{m,n}$ s admit moments of every order, that  $F_\varepsilon^{2+\delta}$  is integrable for  $\delta$  that is small enough and that

$$\frac{1}{n} \sum_{m=0}^{n-1} h_{v_1; v_0}(m/n) h_{v_1; v_0}(m/n) \rightarrow \int_0^1 h_{v_1; v_0} h_{v_2; v_0},$$

using Lemma 10 and the standard monotonous approach for  $\frac{1}{n} \sum_{m=0}^{n-1} h_{v_1; v_0}^2(m/n)$ .

We can then apply the Theorem 6.1 of Ziegler (1997) to prove the claim.  $\square$

**Lemma 12.** *Let  $2/5 < \varepsilon < 1/2$ . The sequence*

$$(v, \alpha) \in N_\varepsilon \times A \mapsto \sqrt{n} \left( \frac{Z^\top R_{v, \alpha}^{-1} Z}{n^{1+2(v-v_0)} \int_0^1 h_{v; v_0}} - 1 \right)$$

of processes converges weakly to

$$\text{GP} \left( 0, (v_1, v_2) \mapsto \frac{\int_0^1 h_{v_1; v_0} h_{v_2; v_0}}{\int_0^1 h_{v_1; v_0} \int_0^1 h_{v_2; v_0}} \right),$$

in  $L^\infty(N_\varepsilon \times A)$ .

*Proof.*

$$\begin{aligned} &\sqrt{n} \left( \frac{Z^\top R_{v, \alpha}^{-1} Z}{n^{1+2(v-v_0)}} - \int_0^1 h_{v; v_0}(x) dx \right) \\ &= \sqrt{n} U_{0,n}^2 \left( \frac{\lambda_{0,n}^0}{n^{1+2(v-v_0)} \lambda_{0,n}} \right) + \frac{1}{\sqrt{n}} \sum_{m=1}^{n-1} U_{m,n}^2 \left( \frac{\lambda_{m,n}^0}{n^{2(v-v_0)} \lambda_{m,n}} - h_{v; v_0}(m/n) \right) \end{aligned}$$



$$+ \frac{1}{\sqrt{n}} \sum_{m=1}^{n-1} B_{m,n} h_{v;v_0}(m/n) + \sqrt{n} \left( \frac{1}{n} \sum_{m=1}^{n-1} h_{v;v_0}(m/n) - \int_0^1 h_{v;v_0}(x) dx \right),$$

with  $B_{m,n} = U_{m,n}^2 - 1$ . Then a classical Borel-Cantelli argument (see, e.g., [Ying, 1991](#), Lemma 4) shows that  $\sup_{0 \leq m \leq n-1} |U_{m,n}^2| = o_P(n^\delta)$  for every  $\delta > 0$ , so

$$\sup_{v \in N_\varepsilon, \alpha \in A} \left| \sqrt{n} U_{0,n}^2 \left( \frac{\lambda_{0,n}^0}{n^{1+2(v-v_0)} \lambda_{0,n}} \right) \right| = o_P(1),$$

and

$$\sup_{v \in N_\varepsilon, \alpha \in A} \left| \frac{1}{\sqrt{n}} \sum_{m=1}^{n-1} U_{m,n}^2 \left( \frac{\lambda_{m,n}^0}{n^{2(v-v_0)} \lambda_{m,n}} - h_{v;v_0}(m/n) \right) \right| = o_P(1).$$

thanks to Lemma 8. Finally Lemma 6, Lemma 11, Slutsky's lemma, and the continuous mapping theorem give the claim observing that the mapping  $v \in N_\varepsilon \mapsto \int_0^1 h_{v;v_0}$  is bounded away from zero.  $\square$

**Lemma 13.** *Let  $v \in (v_0 - 1/2, b_v]$ , we have*

$$g_n(v) = \frac{1}{n} \sum_{m=1}^{n-1} B_{m,n} h_{v;v_0} \left( \frac{m}{n} \right) = o_P(1).$$

*Proof.* Let  $\delta = v - v_0 + 1/2 > 0$ . Then, the equivalence (23) gives

$$\mathbb{E}(g_n^2(v)) = \frac{\text{Var}(B_{0,n})}{n^2} \sum_{m=1}^{n-1} h_{v;v_0}^2 \left( \frac{m}{n} \right) \lesssim \frac{1}{n^2} \sum_{m=1}^l \left( \frac{m}{n} \right)^{4\delta-2} = \frac{1}{n^{4\delta}} \sum_{m=1}^l m^{4\delta-2},$$

which converges to zero no matter how  $\delta$  compares with  $1/4$ .  $\square$

**Lemma 14.** *Let  $0 < \varepsilon < 1/2$ , we have*

$$\sup_{v \in N_\varepsilon, \alpha \in A} \left| \frac{Z^\top R_{v,\alpha}^{-1} Z}{n^{1+2(v-v_0)}} - \int_0^1 h_{v;v_0} \right| = o_P(1).$$

*Proof.* First we have

$$\sup_{v \in N_\varepsilon} \left| \frac{1}{n} \sum_{m=1}^{n-1} B_{m,n} h_{v;v_0} \left( \frac{m}{n} \right) \right| = o_P(1),$$

thanks to Lemma 13, Lemma 15, and ([Van Der Vaart and Wellner, 1996](#), Theorem 1.5.4 and Theorem 1.5.7). Then, proceed as for Lemma 12.  $\square$

**Lemma 15.** *Let  $0 < \varepsilon < 1/2$  and define*

$$g_n : v \in N_\varepsilon \mapsto \frac{1}{n} \sum_{m=1}^{n-1} B_{m,n} h_{v;v_0} \left( \frac{m}{n} \right).$$

*The sequence  $(g_n)$  is asymptotically uniformly equicontinuous in probability for  $(x, y) \mapsto |x - y|$  ([Van Der Vaart and Wellner, 1996](#), Chapter 1.5).*

*Proof.* Let  $1 \leq m \leq \lfloor n/2 \rfloor$  and define

$$\eta_{m,n} : \mathbf{v} \in N_\varepsilon \mapsto \gamma_{2v+1}(m/n) = \sum_{j \in \mathbb{Z}} \frac{1}{|j + m/n|^{2v+1}}.$$

These functions are  $C^1$  and some calculations show that  $(n/m)^{2v+1} \lesssim \eta_{m,n}(\mathbf{v}) \lesssim (n/m)^{2v+1}$  and that  $(n/m)^{2v+1} \ln(n/m) \lesssim \frac{d\eta_{m,n}}{d\mathbf{v}}(\mathbf{v}) \lesssim (n/m)^{2v+1} \ln(n/m)$ , for  $\mathbf{v} \in N_\varepsilon$ ,  $1 \leq m \leq \lfloor n/2 \rfloor$ , and  $n$ .

Write  $\kappa_{m,n} : \mathbf{v} \in N_\varepsilon \mapsto h_{\mathbf{v};v_0}(m/n)$ . Then, for all  $\delta > 0$  and  $\mathbf{v} \in N_\varepsilon$ , we have

$$\left| \frac{d\kappa_{m,n}}{d\mathbf{v}}(\mathbf{v}) \right| \lesssim \left( \frac{n}{m} \right)^{1-2\varepsilon+2\delta}.$$

Now let  $\mathbf{v}_1, \mathbf{v}_2 \in N_\varepsilon$ . If one choose  $p > 1$  and  $\delta > 0$  such that  $p(1 - 2\varepsilon + 2\delta) < 1$ , then we have by Hölder's inequality with  $1/q + 1/p = 1$

$$\begin{aligned} & |g_n(\mathbf{v}_1) - g_n(\mathbf{v}_2)| \\ & \leq \left( \frac{1}{n} \sum_{m=1}^{n-1} B_{m,n}^q \right)^{1/q} \cdot \left( \frac{1}{n} \sum_{m=1}^{n-1} \sup_{\mathbf{v} \in N_\varepsilon} \left| \frac{d\kappa_{m,n}}{d\mathbf{v}}(\mathbf{v}) \right|^p \right)^{1/p} \cdot |\mathbf{v}_1 - \mathbf{v}_2|, \end{aligned}$$

which gives the result since the  $B_{m,n}$ s admit moments of every order.  $\square$

**Lemma 16.** *Let  $2/5 < \varepsilon < 1/2$ . Then, the sequence*

$$(\mathbf{v}, \alpha) \in N_\varepsilon \times A \mapsto \sqrt{n} \left( \mathbb{M}_n(\mathbf{v}, \alpha) - \int_0^1 g_v - \ln \left( \int_0^1 h_{\mathbf{v};v_0} \right) \right)$$

*of processes converges weakly to*

$$\text{GP} \left( 0, (\mathbf{v}_1, \mathbf{v}_2) \mapsto \frac{\int_0^1 h_{\mathbf{v}_1;v_0} h_{\mathbf{v}_2;v_0}}{\int_0^1 h_{\mathbf{v}_1;v_0} \int_0^1 h_{\mathbf{v}_2;v_0}} \right)$$

*in  $L^\infty(N_\varepsilon \times A)$ .*

*Proof.* Let  $\mathbb{E}$  be the set of continuous real-valued functions on the compact  $N_\varepsilon \times A$  and  $\mathbb{D}_\psi \subset \mathbb{E}$  be the subset of strictly positive functions, both endowed with the supremum norm. The mapping  $\psi : g \in \mathbb{D}_\psi \subset \mathbb{E} \mapsto \ln \circ g \in \mathbb{E}$  is Fréchet-differentiable at the unit function and  $\psi'(1) : h \in \mathbb{E} \rightarrow h$ . Then, Theorem 3.9.4 from [Van Der Vaart and Wellner \(1996\)](#) and Lemma 12 show that

$$\sup_{(\mathbf{v}, \alpha) \in N_\varepsilon \times A} \left| \sqrt{n} \ln \left( \frac{Z^\top R_{\mathbf{v}, \alpha}^{-1} Z}{n^{1+2(v-v_0)} \int_0^1 h_{\mathbf{v};v_0}} \right) - \sqrt{n} \left( \frac{Z^\top R_{\mathbf{v}, \alpha}^{-1} Z}{n^{1+2(v-v_0)} \int_0^1 h_{\mathbf{v};v_0}} - 1 \right) \right|$$

is  $o_P(1)$ . Lemma 4 and Slutsky's Lemma allows then to conclude.  $\square$

**Lemma 17.** *The convergence  $\widehat{v}_0 \rightarrow v_0$  holds in probability.*

*Proof.* First for  $\mathbf{v} \in N$  and  $\alpha \in A$ , we have

$$\mathbb{M}_n(\mathbf{v}, \alpha) = \ln(\det(R_{\mathbf{v}, \alpha})) / n + \ln \left( \frac{Z^\top R_{\mathbf{v}, \alpha}^{-1} Z}{n^{1-2v_0}} \right)$$

$$\begin{aligned}
&= \int_0^1 g_v(x) dx + \mathcal{O}(\ln(n)/n) + \ln\left(\frac{Z^\top R_{v,\alpha}^{-1} Z}{n^{1+2(v-v_0)}}\right) \\
&= \mathcal{O}(1) + \ln\left(\frac{Z^\top R_{v,\alpha}^{-1} Z}{n^{1+2(v-v_0)}}\right)
\end{aligned}$$

uniformly thanks to Lemma 4 and by observing that the mapping  $v \in [a_v, b_v] \mapsto \int_0^1 g_v$  is continuous. Now let  $v \in [a_v, v_0 - 1/2 + \varepsilon]$  and  $\alpha \in A$ , Lemma 2 yields

$$\begin{aligned}
\frac{Z^\top R_{v,\alpha}^{-1} Z}{n^{1+2(v-v_0)}} &\gtrsim \frac{1}{n} \sum_{m=1}^l U_{m,n}^2 \left(\frac{m}{n}\right)^{2(v-v_0)} \gtrsim \frac{1}{n} \sum_{m=1}^l U_{m,n}^2 \left(\frac{m}{n}\right)^{-1+2\varepsilon} \\
&= \frac{1}{n^{2\varepsilon}} \sum_{m=1}^l U_{m,n}^2 m^{-1+2\varepsilon} \rightarrow \frac{1}{2^{1+2\varepsilon} \varepsilon}
\end{aligned}$$

in probability using a similar argument as the one from Lemma 13. Considering  $\mathbb{U}_n$  from Section A.1, Lemma 16 gives  $\mathbb{U}_n(v_0) \rightarrow \int_0^1 g_{v_0}$  in probability, so we have

$$\begin{aligned}
&\inf_{v \in [a_v, v_0 - 1/2 + \varepsilon]} \mathbb{U}_n(v) - \mathbb{U}_n(v_0) \\
&\geq C + \inf_{v \in [a_v, v_0 - 1/2 + \varepsilon], \alpha \in A} \ln\left(\frac{Z^\top R_{v,\alpha}^{-1} Z}{n^{1+2(v-v_0)}}\right) - \mathbb{U}_n(v_0) \rightarrow C - \ln(2^{1+2\varepsilon} \varepsilon) - \int_0^1 g_{v_0},
\end{aligned}$$

with the later convergence holding in probability, for a universal constant  $C$ . Letting  $\varepsilon \rightarrow 0$  shows that the above limit can be made arbitrarily high.

Finally let  $0 < \varepsilon < 1/2$ , we have

$$\begin{aligned}
&\sup_{v \in N_\varepsilon} \left| \mathbb{U}_n(v) - \int_0^1 g_v(x) dx - \ln\left(\int_0^1 h_{v;v_0}\right) \right| \\
&\leq \sup_{v \in N_\varepsilon, \alpha \in A} \left| \mathbb{M}_n(v, \alpha) - \int_0^1 g_v(x) dx - \ln\left(\int_0^1 h_{v;v_0}\right) \right| = o_P(1),
\end{aligned}$$

thanks to Lemma 4, Lemma 14 and the continuous mapping theorem applied from the space of functions that are bounded away from zero and infinity to the space of bounded functions. Moreover the function

$$\mathbb{U} : v \in N_\varepsilon \mapsto \int_0^1 g_v(x) dx + \ln\left(\int_0^1 h_{v;v_0}\right)$$

is continuous, and strictly maximized by taking  $v = v_0$  thanks to Jensen inequality and the fact that  $h_{v;v_0}$  is constant only if  $v = v_0$ .  $\square$

**Lemma 18.** *Let  $0 < \beta < 1/4$ . The bound  $\widehat{v}_n - v_0 = o_P(n^{-\beta})$  holds in probability.*

*Proof.* Let  $2/5 < \varepsilon < 1/2$ ,  $0 < \beta < 1/2$  and let us reuse the function  $\mathbb{U}$  from the proof of Lemma 17. We have

$$\mathbb{M}_n(v, \alpha) - \mathbb{U}(v) = \mathcal{O}\left(\frac{\ln(n)}{n}\right) + \ln\left(\frac{Z^\top R_{v,\alpha}^{-1} Z}{n^{1+2(v-v_0)}}\right) - \ln\left(\int_0^1 h_{v;v_0}\right), (v, \alpha) \in N_\varepsilon \times A.$$

So this yields

$$n^\beta \sup_{\mathbf{v} \in N_\varepsilon} |\mathbb{U}_n(\mathbf{v}) - \mathbb{U}(\mathbf{v})| \leq n^\beta \sup_{\mathbf{v} \in N_\varepsilon, \alpha \in A} |\mathbb{M}_n(\mathbf{v}, \alpha) - \mathbb{U}(\mathbf{v})|,$$

which converges in probability to zero thanks to Lemma 16 and Slutsky's Lemma. Moreover,  $\mathbb{U}$  is  $C^3$  and some calculations show that

$$\mathbb{U}''(\mathbf{v}_0) = \int_0^1 \left( \frac{\tilde{\gamma}_{2\mathbf{v}_0+1}}{\gamma_{2\mathbf{v}_0+1}} \right)^2 - \left( \int_0^1 \frac{\tilde{\gamma}_{2\mathbf{v}_0+1}}{\gamma_{2\mathbf{v}_0+1}} \right)^2 > 0,$$

with  $\tilde{\gamma}$  the derivative of  $(\mathbf{v}, x) \mapsto \gamma_{2\mathbf{v}+1}(x)$  with respect to  $\mathbf{v}$ . Then, Lemma 17 and the fact that the maximum of  $\mathbb{U}$  on the compact  $N_\varepsilon$  is unique give the rate  $n^{-\beta/2}$  thanks to a standard Taylor expansion around  $\mathbf{v}_0$ .  $\square$

*Proof of Theorem 1.* Let  $0 < \beta < 1/4$ , Lemma 1 gives

$$\ln(\hat{\phi}_n) = \ln(\phi_0) + \ln\left(\frac{Z^\top R_{\hat{\mathbf{v}}_n, \hat{\alpha}_n}^{-1} Z}{n^{1+2(\hat{\mathbf{v}}_n - \mathbf{v}_0)}}\right) + 2(\hat{\mathbf{v}}_n - \mathbf{v}_0) \ln(n).$$

So

$$\begin{aligned} \frac{n^\beta}{\ln(n)} \left( \ln(\hat{\phi}_n) - \ln(\phi_0) \right) &= \frac{n^\beta}{\ln(n)} \ln\left(\frac{Z^\top K_{\hat{\mathbf{v}}_n, \hat{\alpha}_n}^{-1} Z}{n^{1+2(\hat{\mathbf{v}}_n - \mathbf{v}_0)}}\right) + 2n^\beta (\hat{\mathbf{v}}_n - \mathbf{v}_0) \\ &= \frac{n^\beta}{\ln(n)} \ln\left(\int_0^1 h_{\hat{\mathbf{v}}_n, \mathbf{v}_0}\right) + \frac{n^\beta}{\ln(n)} \left( \ln\left(\frac{Z^\top K_{\hat{\mathbf{v}}_n, \hat{\alpha}_n}^{-1} Z}{n^{1+2(\hat{\mathbf{v}}_n - \mathbf{v}_0)}}\right) - \ln\left(\int_0^1 h_{\hat{\mathbf{v}}_n, \mathbf{v}_0}\right) \right) + 2n^\beta (\hat{\mathbf{v}}_n - \mathbf{v}_0). \end{aligned}$$

The latter converges to zero in probability thanks to Lemma 16, Slutsky's Lemma, Lemma 18, and to the univariate Delta method, observing that the mapping  $\mathbf{v} \mapsto \ln\left(\int_0^1 h_{\mathbf{v}, \mathbf{v}_0}\right)$  is  $C^1$  in a neighborhood of  $\mathbf{v}_0$ .  $\square$

## A.5 Proof of Theorem 2

We start by providing a few cumbersome expressions. Remember that the quantities  $\lambda_{m,n}$  depends on  $\mathbf{v}$  and  $\alpha$  but have their argument dropped to avoid cumbersome notations. The function  $\mathbb{L}_n$  is  $C^3$  for any realization, and

$$\begin{aligned} \mathbb{L}_n(\mathbf{v}, \phi, \alpha) &= \ln(\phi) + \frac{1}{n} \sum_{m=0}^{n-1} \ln(\lambda_{m,n}) + \frac{\phi_0}{n\phi} \sum_{m=0}^{n-1} \frac{\lambda_{m,n}^0 U_{m,n}^2}{\lambda_{m,n}} \\ \frac{\partial \mathbb{L}_n}{\partial \mathbf{v}}(\mathbf{v}, \phi, \alpha) &= \frac{1}{n} \sum_{m=0}^{n-1} \frac{\partial \lambda_{m,n}}{\partial \mathbf{v}} \frac{1}{\lambda_{m,n}} - \frac{\phi_0}{n\phi} \sum_{m=0}^{n-1} \frac{\lambda_{m,n}^0 \frac{\partial \lambda_{m,n}}{\partial \mathbf{v}} U_{m,n}^2}{\lambda_{m,n}^2} \\ \frac{\partial^2 \mathbb{L}_n}{\partial \mathbf{v}^2}(\mathbf{v}, \phi, \alpha) &= \frac{1}{n} \sum_{m=0}^{n-1} \frac{\frac{\partial^2 \lambda_{m,n}}{\partial \mathbf{v}^2} \lambda_{m,n} - \left(\frac{\partial \lambda_{m,n}}{\partial \mathbf{v}}\right)^2}{\lambda_{m,n}^2} \\ &\quad - \frac{\phi_0}{n\phi} \sum_{m=0}^{n-1} \frac{\lambda_{m,n}^0 \left(\frac{\partial^2 \lambda_{m,n}}{\partial \mathbf{v}^2} \lambda_{m,n} - 2\left(\frac{\partial \lambda_{m,n}}{\partial \mathbf{v}}\right)^2\right) U_{m,n}^2}{\lambda_{m,n}^3} \end{aligned}$$

$$\begin{aligned}
\frac{\partial^3 \mathbb{L}_n}{\partial v^3}(v, \phi, \alpha) &= \frac{1}{n} \sum_{m=0}^{n-1} \lambda_{m,n}^{-3} \left( \frac{\partial^3 \lambda_{m,n}}{\partial v^3} \lambda_{m,n} + \frac{\partial^2 \lambda_{m,n}}{\partial v^2} \frac{\partial \lambda_{m,n}}{\partial v} \right. \\
&\quad \left. - 2 \frac{\partial^2 \lambda_{m,n}}{\partial v^2} \frac{\partial \lambda_{m,n}}{\partial v} \right) \lambda_{m,n} - 2 \frac{\partial \lambda_{m,n}}{\partial v} \left( \frac{\partial^2 \lambda_{m,n}}{\partial v^2} \lambda_{m,n} - \left( \frac{\partial \lambda_{m,n}}{\partial v} \right)^2 \right) \\
&\quad - \frac{\phi_0}{n\phi} \sum_{m=0}^{n-1} \lambda_{m,n}^{-4} \lambda_{m,n}^0 \left( \lambda_{m,n} \left( \frac{\partial^3 \lambda_{m,n}}{\partial v^3} \lambda_{m,n} + \frac{\partial^2 \lambda_{m,n}}{\partial v^2} \frac{\partial \lambda_{m,n}}{\partial v} - 4 \frac{\partial^2 \lambda_{m,n}}{\partial v^2} \frac{\partial \lambda_{m,n}}{\partial v} \right) \right. \\
&\quad \left. - 3 \frac{\partial \lambda_{m,n}}{\partial v} \left( \frac{\partial^2 \lambda_{m,n}}{\partial v^2} \lambda_{m,n} - 2 \left( \frac{\partial \lambda_{m,n}}{\partial v} \right)^2 \right) \right) U_{m,n}^2
\end{aligned}$$

Lemma 2 and the following will help to analyze roughly the previous expressions. Exceptionally, the arguments of the  $\lambda_{m,n}$ s are not dropped.

**Lemma 19.** Let  $\delta > 0$ ,  $0 \leq m \leq \lfloor n/2 \rfloor$ ,  $v \in N$ ,  $\alpha \in A$  and  $k \in \{1, 2, 3\}$ . The functions  $\lambda_{m,n}$  are  $C^3$  and:

$$\frac{1}{n} \left| \frac{\partial^k \lambda_{m,n}}{\partial v^k}(v, \alpha) \right| \lesssim \frac{1}{m^{2v+1-\delta}}, \quad \text{if } 1 \leq m \leq \lfloor n/2 \rfloor$$

and

$$1 \lesssim \frac{1}{n} \left| \frac{\partial^k \lambda_{0,n}}{\partial v^k}(v, \alpha) \right| \lesssim 1.$$

*Proof.* We have

$$\left| \frac{\partial^k \lambda_{m,n}}{\partial v^k}(v, \alpha) \right| = \sum_{j \in \mathbb{Z}} \frac{|\ln^k(\alpha^2 + (m+jn)^2)|}{(\alpha^2 + (m+jn)^2)^{v+1/2}} \lesssim \sum_{j \in \mathbb{Z}} \frac{1}{(\alpha^2 + (m+jn)^2)^{v+1/2-\delta/2}}.$$

Lemma 2 gives the result.  $\square$

*Proof of Theorem 2.* Remember that  $A = \{\alpha_0\}$  and consider  $\mathbb{L}_n$  as a function of  $(v, \phi)$  only. We start by elucidating the behavior of the score function.

Let  $e_2(v) = E(\psi_v^2(U))$  for  $v > 0$ . Stein (1999, proof of Theorem 1, Section 6.7) shows that

$$\text{Cov} \left( \frac{\sqrt{n}}{2} \nabla \mathbb{L}_n(v_0, \phi_0) \right) = \underbrace{\begin{pmatrix} 2 \ln^2(n) + 4 \ln(n) e(v_0) + 2 e_2(v_0) & -\frac{\ln(n)}{\phi_0} - \frac{e(v_0)}{\phi_0} \\ -\frac{\ln(n)}{\phi_0} - \frac{e(v_0)}{\phi_0} & \frac{1}{2\phi_0^2} \end{pmatrix}}_{C_n} + \mathcal{O}(n^{-\varepsilon}),$$

for some  $\varepsilon > 0$ .

Define

$$A_n = \frac{2\phi_0}{\sqrt{V(v_0)}} \begin{pmatrix} \frac{1}{2\phi_0} & 0 \\ \ln(n) + e(v_0) & \sqrt{V(v_0)} \end{pmatrix}.$$

One has  $A_n^\top C_n A_n = 2I_2$  and so

$$\text{Cov} \left( \frac{\sqrt{n}}{2} A_n^\top \nabla \mathbb{L}_n(v_0, \phi_0) \right) = A_n^\top \text{Cov} \left( \frac{\sqrt{n}}{2} \nabla \mathbb{L}_n(v_0, \phi_0) \right) A_n \rightarrow 2I_2.$$

Let us show the Lyapunov condition. To do so, write

$$\mathbb{L}_{m,n}(\mathbf{v}, \phi) = \ln(\phi) + \ln(\lambda_{m,n}) + \frac{\phi_0 \lambda_{m,n}^0 U_{m,n}^2}{\phi \lambda_{m,n}},$$

so  $\mathbb{L}_n(\mathbf{v}, \phi) = n^{-1} \sum_{m=0}^{n-1} \mathbb{L}_{m,n}(\mathbf{v}, \phi)$ . Let  $\delta > 0$ , we have

$$\sum_{m=0}^{n-1} \mathbb{E} \left( \left| \frac{\partial \mathbb{L}_{m,n}}{\partial \phi}(\mathbf{v}_0, \phi_0) \right|^{2+\delta} \right) = \frac{1}{\phi_0^{2+\delta} n^{2+\delta}} \sum_{m=0}^{n-1} \mathbb{E} \left( |1 - U_{m,n}^2|^{2+\delta} \right) = \frac{\mathbb{E} \left( |1 - U_{0,1}^2|^{2+\delta} \right)}{\phi_0^{2+\delta} n^{1+\delta}},$$

and

$$\sum_{m=0}^{n-1} \mathbb{E} \left( \left| \frac{\partial \mathbb{L}_{m,n}}{\partial \mathbf{v}}(\mathbf{v}_0, \phi_0) \right|^{2+\delta} \right) = \frac{\mathbb{E} \left( |1 - U_{0,1}^2|^{2+\delta} \right)}{n^{2+\delta}} \sum_{m=0}^{n-1} \left( \frac{\partial \lambda_{m,n}^0}{\lambda_{m,n}^0} \right)^{2+\delta} \lesssim \frac{1}{n^{1+3\delta/4}},$$

using Lemma 2, Lemma 19, and the derivatives expressions. Therefore,

$$\sum_{m=0}^{n-1} \mathbb{E} \left( \|\sqrt{n} A_n^\top \nabla \mathbb{L}_{m,n}(\mathbf{v}_0, \phi_0)\|^{2+\delta} \right) \lesssim n^{1+\delta/2} \ln(n)^{2+\delta} \sum_{m=0}^{n-1} \mathbb{E} \left( \|\nabla \mathbb{L}_{m,n}(\mathbf{v}_0, \phi_0)\|^{2+\delta} \right) \rightarrow 0.$$

The Lyapunov condition is fulfilled, and it is straightforward to verify that the score function is centered, yielding

$$\frac{\sqrt{n}}{2\sqrt{2}} A_n^\top \nabla \mathbb{L}_n(\mathbf{v}_0, \phi_0) \rightsquigarrow \mathcal{N}(0, I_2).$$

Consider now the Hessian matrix. One has

$$\mathbb{E}(\nabla^2 \mathbb{L}_n(\mathbf{v}_0, \phi_0)) = \frac{n}{2} \text{Cov}(\nabla \mathbb{L}_n(\mathbf{v}_0, \phi_0)),$$

so

$$\frac{1}{2} \mathbb{E}(\nabla^2 \mathbb{L}_n(\mathbf{v}_0, \phi_0)) - C_n = \mathcal{O}(n^{-\varepsilon}).$$

Furthermore, using Lemma 2, Lemma 19, and the previous expressions, it is straightforward to check that

$$\text{Var} \left( \frac{\partial^2 \mathbb{L}_n}{(\partial \mathbf{v})^p (\partial \phi)^{2-p}}(\mathbf{v}_0, \phi_0) \right) = \mathcal{O}(n^{-2\varepsilon}),$$

for  $\varepsilon > 0$  small enough, so  $\nabla^2 \mathbb{L}_n(\mathbf{v}_0, \phi_0)/2 - C_n = \mathcal{O}_P(n^{-\varepsilon})$  for some  $\varepsilon > 0$  and therefore  $A_n^\top \nabla^2 \mathbb{L}_n(\mathbf{v}_0, \phi_0) A_n \rightarrow 4I_2$ , in probability.

Let us now roughly bound the third derivatives. For  $\varepsilon > 0$ ,  $m \geq 1$ ,  $k \in \{0, 1, 2, 3\}$ , and  $|\mathbf{v} - \mathbf{v}_0| \leq \varepsilon$ , one has

$$\left| \frac{\partial^k \lambda_{m,n}}{\partial \mathbf{v}^k}(\mathbf{v}_0 + \varepsilon, 1) \right| \leq \left| \frac{\partial^k \lambda_{m,n}}{\partial \mathbf{v}^k}(\mathbf{v}, 1) \right| \leq \left| \frac{\partial^k \lambda_{m,n}}{\partial \mathbf{v}^k}(\mathbf{v}_0 - \varepsilon, 1) \right|.$$

Therefore, using again the previous expressions, Lemma 2, and Lemma 19, it is straightforward to show that

$$\mathbb{E} \left( \sup_{p \in \{0, 1, 2, 3\}, |\mathbf{v} - \mathbf{v}_0| \leq \varepsilon, |\phi - \phi_0| \leq \varepsilon} \left| \frac{\partial^3 \mathbb{L}_n}{(\partial \mathbf{v})^p (\partial \phi)^{3-p}}(\mathbf{v}, \phi) \right| \right) = \mathcal{O}(n^{17\varepsilon}). \quad (25)$$

Now we have

$$0 = \Delta \mathbb{L}_n(\mathbf{v}_0, \phi_0) + \Delta^2 \mathbb{L}_n(\mathbf{v}_0, \phi_0) \begin{pmatrix} \widehat{\mathbf{v}}_n - \mathbf{v}_0 \\ \widehat{\phi}_n - \phi_0 \end{pmatrix} + \mathcal{O}_P \left( n^{17\varepsilon} \left\| \begin{pmatrix} \widehat{\mathbf{v}}_n - \mathbf{v}_0 \\ \widehat{\phi}_n - \phi_0 \end{pmatrix} \right\|^2 \right),$$

thanks to (25) and using Theorem 1 leads to

$$0 = A_n^\top \Delta \mathbb{L}_n(\mathbf{v}_0, \phi_0) + (A_n^\top \Delta^2 \mathbb{L}_n(\mathbf{v}_0, \phi_0) A_n + o_P(1)) A_n^{-1} \begin{pmatrix} \widehat{\mathbf{v}}_n - \mathbf{v}_0 \\ \widehat{\phi}_n - \phi_0 \end{pmatrix},$$

and

$$\sqrt{2n} A_n^{-1} \begin{pmatrix} \widehat{\mathbf{v}}_n - \mathbf{v}_0 \\ \widehat{\phi}_n - \phi_0 \end{pmatrix} = - (A_n^\top \Delta^2 \mathbb{L}_n(\mathbf{v}_0, \phi_0) A_n + o_P(1))^{-1} A_n^\top \Delta \mathbb{L}_n(\mathbf{v}_0, \phi_0) \rightsquigarrow \mathcal{N}(0, I_2),$$

thanks to Slutsky's Lemma. This gives the result.  $\square$

## A.6 Proofs of Theorem 3 and Theorem 4

The posterior mean and the error of the GP does not depend on  $\phi$ , so all derivations will be written with  $\phi = 1$ . We also suppose that  $\phi_0 = 1$  without loss of generality. Furthermore, we will write  $\omega = (\mathbf{v}, \alpha)$  and use the coefficients  $c_j(\omega) = c_j(\mathbf{v}, \alpha)$  defined in Section A.1 to avoid cumbersome expressions in this section.

The sample paths are in  $L^2_{[0,1]}$  almost surely so the convergence also holds with probability one in this sense and the proofs will rely on using Parseval's identity. Note that we assume  $\mathbf{v}_0 > 1/2$  to avoid dealing with conditionally convergent series. Indeed, in this case it is straightforward to check that the expansion (6) converges almost surely absolutely pointwise, so the hypothesis of Proposition 1 are fulfilled.

Let  $\alpha > 0$ ,  $\mathbf{v} > 0$  and  $j \in \mathbb{Z}$ , we have

$$\begin{aligned} 2 \left| c_j(\xi - \widehat{\xi}_n) \right|^2 &= \left( \frac{c_j(\omega) \sum_{j_1 \in \mathbb{Z} \setminus \{0\}} \sqrt{c_{j+j_1}(\omega_0)} U_{1,|j+j_1|n}}{\sum_{j_1 \in \mathbb{Z}} c_{j+nj_1}(\omega)} \right. \\ &\quad \left. - \frac{\sqrt{c_j(\omega_0)} U_{1,|j|} \sum_{j_1 \in \mathbb{Z} \setminus \{0\}} c_{j+nj_1}(\omega)}{\sum_{j_1 \in \mathbb{Z}} c_{j+nj_1}(\omega)} \right)^2 \\ &+ \left( \frac{c_j(\omega) \sum_{j_1 \in \mathbb{Z} \setminus \{0\}} \sqrt{c_{j+nj_1}(\omega_0)} U_{2,|j+nj_1|} \text{sign}(j+nj_1)}{\sum_{j_1 \in \mathbb{Z}} c_{j+nj_1}(\omega)} \right. \\ &\quad \left. - \frac{\sqrt{c_j(\omega_0)} U_{2,|j|} \text{sign}(j) \sum_{j_1 \in \mathbb{Z} \setminus \{0\}} c_{j+nj_1}(\omega)}{\sum_{j_1 \in \mathbb{Z}} c_{j+nj_1}(\omega)} \right)^2. \end{aligned} \quad (26)$$

The two terms in (26) are independent and they are also identically distributed if  $m \notin \{0, n/2\}$ .

*Proof of Theorem 3.* Consider  $m \notin \{0, n/2\}$  and the indexes of the form  $m+nj$ , with  $j \in \mathbb{Z}$ . The two terms in (26) are independent and identically distributed, so there exists a  $\chi^2_2$  distributed variable  $A_{m,j,n}$  such that

$$\left| c_{m+jn}(\xi - \widehat{\xi}_n) \right|^2 = a_{m,j,n}(\mathbf{v}) A_{m,j,n} / 2,$$

with

$$a_{m,j,n}(\mathbf{v}) = c_{m+jn}^2(\boldsymbol{\omega}) \frac{\sum_{j_1 \in \mathbb{Z}} c_{m+nj_1}(\boldsymbol{\omega}) - c_{m+jn}(\boldsymbol{\omega})}{\left(\sum_{j_1 \in \mathbb{Z}} c_{m+nj_1}(\boldsymbol{\omega})\right)^2} + c_{m+jn}(\boldsymbol{\omega}) \left(1 - \frac{c_{m+jn}(\boldsymbol{\omega})}{\sum_{j_1 \in \mathbb{Z}} c_{m+nj_1}(\boldsymbol{\omega})}\right)^2, \quad (27)$$

and the dependence of  $a_{m,j,n}$  on (the fixed)  $\alpha$  be removed for readability. Using Lemma 2 gives, for  $1 \leq m \leq l$  and  $j \neq 0$ :

$$a_{m,j,n}(\mathbf{v}) \lesssim (jn)^{-4v-2} m^{4v+2-2v_0-1} + (jn)^{-2v_0-1}, \quad (28)$$

and

$$a_{m,0,n}(\mathbf{v}) \lesssim n^{-2v_0-1} + m^{4v-2v_0+1} n^{-4v-2}, \quad (29)$$

uniformly in  $j$  and  $m$ . So, this yields

$$\sum_{j \in \mathbb{Z}} a_{m,j,n}(\mathbf{v}) \lesssim n^{-2v_0-1} + n^{-4v-2} m^{4v-2v_0+1}. \quad (30)$$

The first two statements then follow from Lemma 20, Lemma 21, the identity

$$\sum_{j \in \mathbb{Z}} \left| c_{m+jn}(\xi - \widehat{\xi}_n) \right|^2 = \sum_{j \in \mathbb{Z}} \left| c_{n-m+jn}(\xi - \widehat{\xi}_n) \right|^2, \quad (31)$$

for every  $0 \leq m \leq n-1$ , and the Fubini-Tonelli theorem.

For the last statement, let  $1 \leq m \leq l$ . We have

$$\begin{aligned} \mathbb{E} \left( \left| c_{m+jn}(\xi - \widehat{\xi}_n) \right|^2 \right) &= c_{m+jn}^2(\boldsymbol{\omega}) \frac{\sum_{j_1 \in \mathbb{Z}} c_{m+nj_1}(\boldsymbol{\omega})}{\left(\sum_{j_1 \in \mathbb{Z}} c_{m+nj_1}(\boldsymbol{\omega})\right)^2} \\ &\quad + c_{m+jn}(\boldsymbol{\omega}) \left(1 - 2 \frac{c_{m+jn}(\boldsymbol{\omega})}{\sum_{j_1 \in \mathbb{Z}} c_{m+nj_1}(\boldsymbol{\omega})}\right), \end{aligned}$$

for every  $j \in \mathbb{Z}$  and therefore

$$\begin{aligned} &\sum_{j \in \mathbb{Z}} \mathbb{E} \left( \left| c_{m+jn}(\xi - \widehat{\xi}_n) \right|^2 \right) \\ &= (1 + \mathcal{O}(m^{-2})) \sum_{j \in \mathbb{Z}} |m+jn|^{-4v-2} \frac{\sum_{j_1 \in \mathbb{Z}} |m+nj_1|^{-2v_0-1}}{\left(\sum_{j_1 \in \mathbb{Z}} |m+nj_1|^{-2v-1}\right)^2} \\ &\quad + |m+jn|^{-2v_0-1} \left(1 - 2 \frac{|m+jn|^{-2v-1}}{\sum_{j_1 \in \mathbb{Z}} |m+nj_1|^{-2v-1}}\right) \\ &= \frac{(1 + \mathcal{O}(m^{-2}))}{n^{2v_0+1}} \vartheta_{\mathbf{v}; \mathbf{v}_0}(m/n), \end{aligned}$$

using Lemma 3 and after a few algebraic manipulations. If  $v \geq (v_0 - 1/2)/2$ , then the function  $\vartheta_{\mathbf{v}; \mathbf{v}_0}$  has a finite limit in zero. If  $(v_0 - 1)/2 < v < (v_0 - 1/2)/2$ , then a taylor



expansion of its derivative shows that it is decreasing in a neighborhood of zero. This shows:

$$\frac{1}{n} \sum_{m=1}^l \vartheta_{\mathbf{v}; \mathbf{v}_0}(m/n) \rightarrow \int_0^{1/2} \vartheta_{\mathbf{v}; \mathbf{v}_0}, \quad (32)$$

and Parseval's identity and the Fubini-Tonelli theorem gives

$$n^{2\nu_0} \mathbb{E}(\text{ISE}_n(\mathbf{v}, \alpha; \xi)) = o(1) + \frac{2}{n} \sum_{m=1}^l (1 + \mathcal{O}(m^{-2})) \vartheta_{\mathbf{v}; \mathbf{v}_0}(m/n) \rightarrow \int_0^1 \vartheta_{\mathbf{v}; \mathbf{v}_0},$$

killing the  $\mathcal{O}(m^{-2})$  term by splitting by how  $m$  compares with  $\sqrt{n}$  as in the proof of Lemma 8.  $\square$

**Lemma 20.** *Let  $n \geq 2$  be even, one has*

$$\mathbb{E} \left( \sup_{\mathbf{v} \in N, \alpha \in A} \sum_{j \in \mathbb{Z}} \left| c_{n/2+jn}(\xi - \widehat{\xi}_n) \right|^2 \right) \lesssim n^{-2\nu_0-1}.$$

*Proof.* First, observe that  $\sum_{j \in \mathbb{Z}} \left| c_{n/2+jn}(\xi - \widehat{\xi}_n) \right|^2 = 2 \sum_{j \geq 0} \left| c_{n/2+jn}(\xi - \widehat{\xi}_n) \right|^2$ .

Let  $\mathbf{v} \in N$ ,  $\alpha \in A$ , and  $j \geq 0$ . One has

$$\begin{aligned} & 2 \left| c_{n/2+jn}(\xi - \widehat{\xi}_n) \right|^2 \left( \sum_{j_1 \in \mathbb{Z}} c_{n/2+nj_1}(\omega) \right)^2 \\ &= \left( \left( c_{n/2+jn}(\omega) \sqrt{c_{n/2+jn}(\omega_0)} - \sqrt{c_{n/2+jn}(\omega_0)} \sum_{j_1 \in \mathbb{Z} \setminus \{j\}} c_{n/2+nj_1}(\omega) \right) U_{1,|n/2+jn|} \right. \\ & \quad \left. + 2c_{n/2+jn}(\omega) \sum_{j_1=0, j_1 \neq |j|}^{+\infty} \sqrt{c_{n/2+j_1n}(\omega_0)} U_{1,|n/2+j_1n|} \right)^2 \\ & \quad + \left( \sqrt{c_{n/2+jn}(\omega_0)} \sum_{j_1 \in \mathbb{Z}} c_{n/2+nj_1}(\omega) U_{2,|n/2+jn|} \right)^2 \end{aligned}$$

So

$$2 \left| c_{n/2+jn}(\xi - \widehat{\xi}_n) \right|^2 = d_{j,n}^2 D_{j,n}^2 + b_{j,n}^2 B_{j,n}^2$$

with  $D_{j,n}$  and  $B_{j,n}$  standard Gaussians and

$$\begin{aligned} d_{j,n}^2 &\leq c_{n/2+jn}(\omega_0) + 4c_{n/2+jn}^2(\omega) \frac{\sum_{j_1=0}^{+\infty} c_{n/2+j_1n}(\omega_0)}{\left( \sum_{j_1 \in \mathbb{Z}} c_{n/2+nj_1}(\omega) \right)^2} \\ &\lesssim ((j+1/2)n)^{-2\nu_0-1} + (j+1/2)^{-4\nu-2} n^{-2\nu_0-1} \end{aligned}$$

uniformly in  $\mathbf{v}$  and  $\alpha$  using Lemma 2 and the inequality  $c_{n/2+jn}(\omega) \leq ((j+1/2)n)^{-2\nu-1}$ . Similarly, one has  $b_{j,n}^2 \lesssim ((j+1/2)n)^{-2\nu_0-1}$ . The statement then follows using the Fubini-Tonelli theorem.  $\square$

**Lemma 21.** *Let  $n \geq 2$  and  $p \geq 1$ . Then, there exists standard Gaussians  $\{A_{j,n}, j \in \mathbb{Z}\}$  and  $\varepsilon > 0$  such that*

$$\sum_{j \in n\mathbb{Z}} \left| c_{jn}(\xi - \widehat{\xi}_n) \right|^{2p} \lesssim A_{0,n}^{2p} (n^{-2\nu_0-1} + n^{-4\nu-2})^p + \sum_{j \in \mathbb{Z} \setminus \{0\}} A_{j,n}^{2p} j^{-1-\varepsilon} (n^{-2\nu_0-1} + n^{-4\nu-2})^p,$$

uniformly in  $\nu \in N$  and  $\alpha \in A$ .

*Proof.* Let  $\nu \in N$  and  $\alpha \in A$ . For  $j \in \mathbb{Z} \setminus \{0\}$ , one has

$$\begin{aligned} & 2 \left| c_{jn}(\xi - \widehat{\xi}_n) \right|^2 \left( \sum_{j_1 \in \mathbb{Z}} c_{nj_1}(\omega) \right)^2 \\ &= \left( \left( c_{jn}(\omega) \sqrt{c_{jn}(\omega_0)} - \sqrt{c_{jn}(\omega_0)} \sum_{j_1 \in \mathbb{Z} \setminus \{j\}} c_{nj_1}(\omega) \right) U_{1,|jn|} \right. \\ & \quad \left. + c_{jn}(\omega) \sqrt{c_0(\omega_0)} U_{1,0} + 2c_{jn}(\omega) \sum_{j_1=1, j_1 \neq j}^{+\infty} \sqrt{c_{j_1 n}(\omega_0)} U_{1,|j_1 n|} \right)^2 \\ & \quad + \left( \sqrt{c_{jn}(\omega_0)} \sum_{j_1 \in \mathbb{Z}} c_{nj_1}(\omega) U_{2,|jn|} \right)^2 \end{aligned}$$

Moreover,

$$\begin{aligned} & 2 \left| c_0(\xi - \widehat{\xi}_n) \right|^2 \left( \sum_{j_1 \in \mathbb{Z}} c_{nj_1}(\omega) \right)^2 = \left( \sqrt{c_0(\omega_0)} \sum_{j_1 \in \mathbb{Z} \setminus \{0\}} c_{nj_1}(\omega) U_{1,|0|} \right. \\ & \quad \left. + 2c_0(\omega) \sum_{j_1=1}^{+\infty} \sqrt{c_{j_1 n}(\omega_0)} U_{1,|j_1 n|} \right)^2 \end{aligned}$$

Then, proceed as in Lemma 20 to conclude, observing that  $c_{nj_1}(\omega) \lesssim (|j_1|n)^{-2\nu-1}$  for  $j_1 \neq 0$ .  $\square$

**Lemma 22.** *Let  $\varepsilon > 0$ . With the notations of Theorem 4, we have:*

$$\mathbb{P}(\widehat{\nu}_n \leq \nu_0 - 1/2 - \varepsilon) \lesssim e^{-C\sqrt{n}},$$

for some  $C > 0$ .

*Proof.* First, one has

$$\begin{aligned} \mathbb{P}(\widehat{\nu}_n \leq \nu_0 - 1/2 - \varepsilon) &\leq \mathbb{P} \left( \inf_{a_\nu \leq \nu \leq \nu_0 - 1/2 - \varepsilon} \mathbb{M}_n(\nu, \alpha) \leq \inf_{\nu_0 - 1/2 - \varepsilon \leq \nu \leq b_\nu} \mathbb{M}_n(\nu, \alpha) \right) \\ &\leq \mathbb{P} \left( \inf_{a_\nu \leq \nu \leq \nu_0 - 1/2 - \varepsilon} \mathbb{M}_n(\nu, \alpha) - \mathbb{M}_n(\nu_0, \alpha) \leq 0 \right). \end{aligned}$$

Then, let  $a_v \leq v \leq v_0 - 1/2 - \varepsilon$ , we have, using Lemma 4, Lemma 2, and Jensen inequality:

$$\begin{aligned}
M_n(v, \alpha) &= \mathcal{O}\left(\frac{\ln(n)}{n}\right) + \ln\left(\frac{Z^T R_{v, \alpha}^{-1} Z}{n}\right) + \int_0^1 g_v + 2(v_0 - v) \ln(n) \\
&= \mathcal{O}(1) + \ln\left(\frac{Z^T R_{v, \alpha}^{-1} Z}{n}\right) + 2(v_0 - v) \ln(n) \\
&\geq \mathcal{O}(1) + \ln\left(\frac{\sum_{m=1}^l U_{m,n}^2 m^{2(v-v_0)}}{n^{1+2(v-v_0)}}\right) \\
&= \mathcal{O}(1) + \ln\left(\frac{1}{n} \sum_{m=1}^l U_{m,n}^2 \left(\frac{m}{n}\right)^{2(v-v_0)}\right) \\
&\geq \mathcal{O}(1) + \ln\left(\frac{1}{n} \sum_{m=1}^l U_{m,n}^2 \left(\frac{m}{n}\right)^{-1-2\varepsilon}\right) \\
&\geq \mathcal{O}(1) + 2\varepsilon \ln(n) + \ln\left(\sum_{m=1}^l U_{m,n}^2 m^{-1-2\varepsilon}\right) \\
&\geq \mathcal{O}(1) + 2\varepsilon \ln(n) + \ln\left(\sum_{m=1}^{\lfloor \sqrt{n} \rfloor} U_{m,n}^2 m^{-1-2\varepsilon}\right) \\
&\geq \mathcal{O}(1) + 2\varepsilon \ln(n) + \ln\left(\sum_{m=1}^{\lfloor \sqrt{n} \rfloor} U_{m,n}^2 \lfloor \sqrt{n} \rfloor^{-1-2\varepsilon}\right) \\
&\geq \mathcal{O}(1) + \varepsilon \ln(n) + \ln\left(\frac{1}{\lfloor \sqrt{n} \rfloor} \sum_{m=1}^{\lfloor \sqrt{n} \rfloor} U_{m,n}^2\right) \\
&\geq \mathcal{O}(1) + \varepsilon \ln(n) + \frac{1}{\lfloor \sqrt{n} \rfloor} \sum_{m=1}^{\lfloor \sqrt{n} \rfloor} \ln(U_{m,n}^2)
\end{aligned}$$

with a uniform big- $\mathcal{O}$ . Let  $\delta > 0$  and  $t > 0$ , we have

$$\begin{aligned}
\mathbb{P}\left(-\frac{1}{\lfloor \sqrt{n} \rfloor} \sum_{m=1}^{\lfloor \sqrt{n} \rfloor} \ln(U_{m,n}^2) \geq \delta\right) &= \mathbb{P}\left(e^{-\frac{t}{\lfloor \sqrt{n} \rfloor} \sum_{m=1}^{\lfloor \sqrt{n} \rfloor} \ln(U_{m,n}^2)} \geq e^{t\delta}\right) \\
&= \mathbb{P}\left(\left(\prod_{m=1}^{\lfloor \sqrt{n} \rfloor} U_{m,n}^{-2}\right)^{\frac{t}{\lfloor \sqrt{n} \rfloor}} \geq e^{t\delta}\right) \\
&= \mathbb{P}\left(\prod_{m=1}^{\lfloor \sqrt{n} \rfloor} |U_{m,n}|^{-2t} \geq e^{t\delta \lfloor \sqrt{n} \rfloor}\right) \\
&\leq e^{-\delta \lfloor \sqrt{n} \rfloor / 4} \mathbb{E}\left(|U_{1,1}|^{-1/2}\right)^{\lfloor \sqrt{n} \rfloor},
\end{aligned}$$

with  $t = 1/4$  and  $C = \mathbb{E}\left(U_{1,1}^{-1/2}\right) < +\infty$ . Therefore:

$$C^{\lfloor \sqrt{n} \rfloor} e^{-\delta \lfloor \sqrt{n} \rfloor / 4} = e^{(\ln(C) - \delta/4) \lfloor \sqrt{n} \rfloor},$$

which converges exponentially to zero if  $\delta$  is high enough.

Furthermore, Lemma 2 gives

$$\mathbb{M}_n(v_0, \alpha) = \mathcal{O}(1) + \ln \left( \frac{Z^\top R_{v_0, \alpha}^{-1} Z}{n} \right) = \mathcal{O}(1) + \ln \left( n^{-1} \sum_{m=0}^{n-1} U_{m,n}^2 \right),$$

and we have

$$\mathbb{P} \left( \ln \left( n^{-1} \sum_{m=0}^{n-1} U_{m,n}^2 \right) \geq \delta \right) = \mathbb{P} \left( n^{-1} \sum_{m=0}^{n-1} U_{m,n}^2 \geq e^\delta \right) \leq e^{-Cn},$$

for some  $C > 0$  if  $\delta > 0$  is high enough, using also a Chernoff bound argument. Now, putting all the pieces together yields:

$$\begin{aligned} & \inf_{a_v \leq v \leq v_0 - 1/2 - \varepsilon} \mathbb{M}_n(v, \alpha) - \mathbb{M}_n(v_0, \alpha) \\ & \geq \mathcal{O}(1) + \varepsilon \ln(l) + \frac{1}{\lfloor \sqrt{n} \rfloor} \sum_{m=1}^{\lfloor \sqrt{n} \rfloor} \ln(U_{m,n}^2) - \ln \left( n^{-1} \sum_{m=0}^{n-1} U_{m,n}^2 \right) \end{aligned}$$

giving the result thanks to the pigeonhole principle.  $\square$

*Proof of Theorem 4.* Let  $\varepsilon > 0$  and use the notations (27). The functions  $a_{m,j,n}$  are  $C^1$  and, for  $v \in [v_0 - 1/2 - \varepsilon, b_v]$ , the derivatives of the  $a_{m,j,n}$ s lead to tedious expressions, which in turns lead, with Lemma 2 and Lemma 19, to

$$\begin{aligned} |a'_{m,0,n}(v)| & \lesssim m^\delta n^{-2v_0-1} + m^{4v-2v_0+1} n^{-4v-2+\delta} \\ & \lesssim m^\delta n^{-2v_0-1} + m^{-2v_0+1} n^{-2+\delta} (m/n)^{2v_0-2+4\varepsilon} \\ & = m^\delta n^{-2v_0-1} + m^{-1+4\varepsilon} n^{-2v_0+\delta-4\varepsilon} \end{aligned}$$

and, for  $j \neq 0$ , to

$$\begin{aligned} |a'_{m,j,n}(v)| & \lesssim (|j|n)^{-4v-2+\delta} m^{-2v_0-1} + (|j|n)^{-2v_0-2v-2+\delta} m^{2v+1} \\ & \lesssim |j|^{-3/2} n^{-2v_0-4\varepsilon+\delta} m^{-2v_0-1} + |j|^{-3/2} n^{-2v_0-2+\delta} m^1 (m/n)^{v_0-1+2\varepsilon} \\ & = |j|^{-3/2} n^{-2v_0-4\varepsilon+\delta} m^{-2v_0-1} + |j|^{-3/2} n^{-3v_0-1+\delta-2\varepsilon} m^{v_0+2\varepsilon} \end{aligned}$$

for  $\delta > 0$  small enough and uniformly in  $1 \leq m \leq l$ ,  $j \neq 0$  and  $v \in [v_0 - 1/2 - \varepsilon, b_v]$ . Then,

$$\begin{aligned} & \sum_{m=1}^l \sum_{j \in \mathbb{Z}} \mathbb{E} \left( A_{m,j,n} |a_{m,j,n}(\widehat{v}_n) - a_{m,j,n}(v_0)| \mathbb{1}_{\widehat{v}_n \geq v_0 - 1/2 - \varepsilon} \right) \\ & \leq \sum_{m=1}^l \sum_{j \in \mathbb{Z}} \mathbb{E} \left( A_{m,j,n} |\widehat{v}_n - v_0| \sup_{v_0 - 1/2 - \varepsilon \leq v \leq b_v} |a'_{m,j,n}(v)| \right) \\ & = \sqrt{\mathbb{E}(A_{1,0,1}^2)} \sqrt{\mathbb{E}((\widehat{v}_n - v_0)^2)} \sum_{m=1}^l \sum_{j \in \mathbb{Z}} \sup_{v_0 - 1/2 - \varepsilon \leq v \leq b_v} |a'_{m,j,n}(v)| = o(n^{-2v_0}), \end{aligned}$$

using the above inequalities and Theorem 1. Therefore, Lemma 20, Lemma 21, the identity (31), and the Fubini-Tonelli theorem shows that

$$\mathbb{E} \left( |\text{ISE}_n(\widehat{v}_n, \alpha; \xi) - \text{ISE}_n(v_0, \alpha; \xi)| \mathbb{1}_{\widehat{v}_n \geq v_0 - 1/2 - \varepsilon} \right) = o(n^{-2v_0}).$$

Furthermore, using again the Fubini-Tonelli theorem yields

$$\begin{aligned}
& \mathbb{E} \left( \sum_{m=1}^l \sum_{j \in \mathbb{Z}} \left| c_{m+jn}(\xi - \widehat{\xi}_n) \right|^2 \mathbb{1}_{\widehat{v}_n \leq v_0 - 1/2 - \varepsilon} \right) \\
&= \sum_{m=1}^l \sum_{j \in \mathbb{Z}} \mathbb{E} \left( a_{m,j,n}(\widehat{v}_n) A_{m,j,n} \mathbb{1}_{\widehat{v}_n \leq v_0 - 1/2 - \varepsilon} \right) \\
&\leq \sum_{m=1}^l \sum_{j \in \mathbb{Z}} \sup_{a_v \leq v \leq v_0 - 1/2 - \varepsilon} a_{m,j,n}(v) \mathbb{E} \left( A_{m,j,n} \mathbb{1}_{\widehat{v}_n \leq v_0 - 1/2 - \varepsilon} \right) \\
&\leq \sqrt{\mathbb{E} \left( A_{1,0,1}^2 \right)} \sqrt{\mathbb{E} \left( \mathbb{1}_{\widehat{v}_n \leq v_0 - 1/2 - \varepsilon} \right)} \sum_{m=1}^l \sum_{j \in \mathbb{Z}} \sup_{a_v \leq v \leq v_0 - 1/2 - \varepsilon} a_{m,j,n}(v) \\
&\leq \sqrt{\mathbb{E} \left( A_{1,0,1}^2 \right)} \sqrt{\mathbb{E} \left( \mathbb{1}_{\widehat{v}_n \leq v_0 - 1/2 - \varepsilon} \right)} n^{-4a_v - 2 + 4\varepsilon} \\
&= o \left( n^{-2v_0} \right),
\end{aligned}$$

using inequalities (28) and (29), and Lemma 22. Then, the case  $m = n/2$  is treated by Lemma 20 and the case  $m = 0$  can be managed similarly using Lemma 21.

Finally, one can treat

$$\mathbb{E} \left( \text{ISE}_n(v_0, \alpha; \xi) \mathbb{1}_{\widehat{v}_n \leq v_0 - 1/2 - \varepsilon} \right)$$

the same way and the desired result follows.  $\square$

## A.7 Proofs of Section 4

*Proof of Proposition 2.* The proof is based on the observation that

$$Z^T R_{v,\alpha}^{-1} Z = \sum_{m=0}^{n-1} \frac{|\sum_{j \in m+n\mathbb{Z}} c_j(f)|^2}{\sum_{j \in m+n\mathbb{Z}} c_j(v, \alpha)}.$$

We give a full proof only for the third assumption. The proof for the second assumption is very similar and the first is a particular case.

The definition of  $v_0(f)$  yields

$$|c_j(f)| \lesssim |j|^{-v_0(f)-1+\delta/2} \quad \text{and} \quad |c_j(f)| \gtrsim |j|^{-v_0(f)-1-\delta} \quad (33)$$

for all  $\delta > 0$ .

Let  $\varepsilon > 0$ ,  $a_v \leq v \leq v_0(f) - \varepsilon$ ,  $\alpha \in A$ , and  $p \in \mathbb{Z}$  such that  $c_p(f) \neq 0$ . Then, Lemma 4 gives

$$\begin{aligned}
\mathbb{M}_n^f(v, \alpha) &= 2(v_0(f) - v) \ln(n) + \mathcal{O}(1) + \ln \left( \sum_{m=0}^{n-1} \frac{|\sum_{j \in m+n\mathbb{Z}} c_j(f)|^2}{\sum_{j \in m+n\mathbb{Z}} c_j(v, \alpha)} \right) \\
&\geq 2\varepsilon \ln(n) + \mathcal{O}(1) + \ln \left( \frac{|\sum_{j \in p+n\mathbb{Z}} c_j(f)|^2}{\sum_{j \in p+n\mathbb{Z}} c_j(v, \alpha)} \right) \\
&= 2\varepsilon \ln(n) + \mathcal{O}(1)
\end{aligned}$$

uniformly by Lemma 2 and since  $|\sum_{j \in p+n\mathbb{Z}} c_j(f)| \rightarrow c_p(f)$  by (33).

Moreover, for any  $\alpha \in A$ , we have:

$$\begin{aligned} \mathbb{M}_n^f(v_0(f) + 1/2, \alpha) &= \mathcal{O}(1) + \ln \left( n^{-1} \sum_{m=0}^{n-1} \frac{|\sum_{j \in m+n\mathbb{Z}} c_j(f)|^2}{\sum_{j \in m+n\mathbb{Z}} c_j(v_0(f) + 1/2, \alpha)} \right) \\ &\leq -\ln(n) + \mathcal{O}(1) + \ln \left( \mathcal{O}(1) + \sum_{m=1}^{n-1} \frac{\sum_{j \in \mathbb{Z}} |m + jn|^{-2v_0(f) - 2 + \varepsilon}}{m^{-2v_0(f) - 2}} \right) \\ &= -\ln(n) + \mathcal{O}(1) + \ln \left( \frac{(n-1)^{1+\varepsilon}}{1+\varepsilon} \right) \\ &= \varepsilon \ln(n) + \mathcal{O}(1) \end{aligned}$$

using Lemma 2 and (33) with  $\delta = \varepsilon$ . This completes the proof.  $\square$

*Proof of Proposition 4.* Let  $v > v_0(f)$  and  $\alpha > 0$ . Then, Lemma 3 and Lemma 4 yield:

$$\begin{aligned} \mathbb{M}_n^f(v, \alpha) &= (2v_0(f) + 1) \ln(n) + n^{-1} \ln(\det(R_{v, \alpha})) + \ln \left( \frac{Z^T R_{v, \alpha}^{-1} Z}{n} \right) \\ &= (2v_0(f) + 1) \ln(n) + \int_0^1 g_v + \mathcal{O} \left( \frac{\ln(n)}{n} \right) + \ln \left( n^{-2v-1} \sum_{m=0}^{n-1} \frac{|\sum_{j \in m+n\mathbb{Z}} c_j(f)|^2}{\sum_{j \in m+n\mathbb{Z}} c_j(v, \alpha)} \right) \\ &= (2v_0(f) + 1) \ln(n) + \int_0^1 g_v + \mathcal{O} \left( \frac{\ln(n)}{n} \right) \\ &+ \ln \left( \mathcal{O}(n^{-2v-1}) + \sum_{m=1}^{n-1} \frac{(1 + \mathcal{O}(m^{-2})) |\sum_{j \in m+n\mathbb{Z}} c_j(f)|^2}{\gamma(2v+1; m/n)} \right) \\ &= \int_0^1 g_v + \mathcal{O} \left( \frac{\ln(n)}{n} \right) \\ &+ \ln \left( \mathcal{O}(n^{2(v_0(f)-v)}) + n^{-1} \sum_{m=1}^{n-1} \frac{(1 + \mathcal{O}(m^{-2})) |n^{v_0(f)+1} \sum_{j \in m+n\mathbb{Z}} c_j(f)|^2}{\gamma(2v+1; m/n)} \right) \\ &= \int_0^1 g_v + \mathcal{O} \left( \frac{\ln(n)}{n} \right) \\ &+ \ln \left( \mathcal{O}(n^{2(v_0(f)-v)}) + \mathcal{O}(n^{v_0(f)-v}) + n^{-1} \sum_{m=1}^{n-1} \frac{\gamma^2(v_0(f)+1; m/n)}{\gamma(2v+1; m/n)} \right), \end{aligned}$$

by proceeding as in the proof of Lemma 8. To conclude, observe that  $\gamma^2(v_0(f)+1; \cdot) / \gamma(2v+1; \cdot)$  is non-increasing in a neighborhood of zero if  $v < v_0(f) + 1/2$  and has a finite limit otherwise.  $\square$

## References

M. L. Stein. *Interpolation of spatial data*. Springer Series in Statistics. Springer-Verlag, New York, 1999. Some theory for Kriging.

- Thomas Santner, Brian Williams, and William Notz. *The Design and Analysis Computer Experiments*. 01 2003.
- C. E. Rasmussen and C. K. I. Williams. *Gaussian Processes for Machine Learning*. Adaptive Computation and Machine Learning. MIT Press, Cambridge, MA, USA, 2006.
- B. Matérn. *Spatial Variation*. Springer-Verlag New York, 1986.
- F. Bachoc. Asymptotic analysis of maximum likelihood estimation of covariance parameters for Gaussian processes: an introduction with proofs. In *Advances in Contemporary Statistics and Econometrics*, pages 283–303. Springer, 2021.
- K. V. Mardia and R. J. Marshall. Maximum likelihood estimation of models for residual covariance in spatial regression. *Biometrika*, 71(1):135–146, 1984.
- F. Bachoc. Asymptotic analysis of the role of spatial sampling for covariance parameter estimation of Gaussian processes. *Journal of Multivariate Analysis*, 125:1–35, 2014.
- Z. Ying. Asymptotic properties of a maximum likelihood estimator with data from a Gaussian process. *Journal of Multivariate analysis*, 36(2):280–296, 1991.
- Z. Ying. Maximum likelihood estimation of parameters under a spatial sampling scheme. *The Annals of Statistics*, pages 1567–1590, 1993.
- A. W. van der Vaart. Maximum likelihood estimation under a spatial sampling scheme. *The Annals of Statistics*, pages 2049 – 2057, 1996.
- H. Zhang. Inconsistent estimation and asymptotically equal interpolations in model-based geostatistics. *Journal of the American Statistical Association*, 99(465):250–261, 2004.
- W.-L. Loh. Fixed-domain asymptotics for a subclass of matérn-type Gaussian random fields. *The Annals of Statistics*, 33, 2006.
- E. Anderes. On the consistent separation of scale and variance for Gaussian random fields. *The Annals of Statistics*, 38(2):870–893, 2010.
- Michael L. Stein. Spline smoothing with an estimated order parameter. *The Annals of Statistics*, 21(3):1522–1544, 1993a.
- Y. Chen, H. Owhadi, and A. Stuart. Consistency of empirical bayes and kernel flow for hierarchical parameter estimation. *Mathematics of Computation*, 90(332):2527–2578, 2021.
- T. Karvonen. Asymptotic bounds for smoothness parameter estimates in gaussian process regression. *arXiv preprint arXiv:2203.05400*, 2022.
- B. Szabó, A. W. van Der Vaart, and J. H. van Zanten. Frequentist coverage of adaptive nonparametric Bayesian credible sets. *The Annals of Statistics*, 43(4):1391–1428, 2015.
- B. T. Knapik, B. T. Szabó, A. W. van Der Vaart, and J. H. van Zanten. Bayes procedures for adaptive inference in inverse problems for the white noise model. *Probability Theory and Related Fields*, 164(3):771–813, 2016.

- T. Karvonen, G. Wynne, F. Tronarp, C. Oates, and S. Sarkka. Maximum likelihood estimation and uncertainty quantification for Gaussian process approximation of deterministic functions. *SIAM/ASA Journal on Uncertainty Quantification*, 8(3):926–958, 2020.
- Michael L. Stein. A simple condition for asymptotic optimality of linear predictions of random fields. *Statistics and Probability Letters*, 17(5):399–404, August 1993b.
- G. Matheron. The theory of regionalized variables and its applications. Technical Report Les cahiers du CMM de Fontainebleau, Fasc. 5, Ecole des Mines de Paris, 1971.
- W. Wang and B.-Y. Jing. Gaussian process regression: Optimality, robustness, and relationship with kernel ridge regression. *Journal of Machine Learning Research*, 23(193):1–67, 2022.
- P. Craven and G. Wahba. Smoothing noisy data with spline functions. Estimating the correct degree of smoothing by the method of generalized cross-validation. *Numerische Mathematik*, 1979.
- P. J. Brockwell and R. A. Davis. *Time series: theory and methods*. Springer science & business media, 1987.
- A. G. Postnikov. *Introduction to analytic number theory*, volume 68. American Mathematical Soc., 1988.
- K. Ziegler. Functional central limit theorems for triangular arrays of function-indexed processes under uniformly integrable entropy conditions. *journal of multivariate analysis*, 62(2):233–272, 1997.
- A. W. Van Der Vaart and J. A. Wellner. *Weak convergence and empirical processes: with applications to statistics*. Springer Science & Business Media, 1996.